

SURFEX OFF-LINE

User's Guide

surfex version : v3

P. Le Moigne

10 janvier 2008

Table des matières

I	How to install the software	7
1	Export off-line version of SURFEX	9
II	Off-line guide	13
2	Using the externalized surface in off-line mode	15
2.1	Case study	15
2.2	The input files	15
3	Namelist options	17
3.1	namelist for PGD	17
3.2	namelist for PREP	18
3.3	namelist for RUN	18
4	Data input files	19
4.1	netcdf format files	19
4.1.1	FORCING.nc	19
4.1.2	PARAMS.nc	20
4.2	ascii format files	20
4.2.1	PGD.txt	22
4.2.2	PREP.txt	23
4.2.3	Extracting 2d fields	23
4.2.4	I/O diagram	25
5	Architecture of the off-line driver	27
5.1	Initialization	27
5.2	Temporal loop	27
5.2.1	coupling surface with the atmosphere	28
5.2.2	writing out prognostic variables	28
5.2.3	compute diagnostics	28

5.2.4	writing out diagnostics variables	28
5.3	writing out restart file	28
6	Some output of off-line simulation	29
6.1	Examples of prognostic variables output	29
6.2	list of available variables	30
III	Users guide	33
7	Overview of the externalized surface sequence	35
7.1	The sequence	35
7.2	The atmospheric models using the externalized surface	35
7.2.1	in offline mode	35
7.2.2	in MESONH	35
7.2.3	in AROME	36
8	The physiographic fields	37
8.1	Overview of physiographic fields computation : PGD	37
8.1.1	Choice of the grid	37
8.1.2	Choice of the physiographic fields	37
8.2	Choice of the surface schemes	46
8.3	Definition of the grid	47
8.3.1	Choice of the grid type	47
8.3.2	Conformal projection grids (Mercator, Lambert, Polar stereographic)	48
8.3.3	Cartesian grids	49
8.3.4	Regular longitude-latitude grids	50
8.3.5	Regular Lambert grids	51
8.3.6	Gaussian grids	52
8.4	Land cover fractions	53
8.5	Orography and subgrid orography	54
8.6	Namelist for ISBA scheme	55
8.7	Namelist to add user's own fields	56
8.8	Namelist for chemistry anthropogenic emissions	57
9	Initialization of the prognostic fields	59
9.1	Overview of fields computation : PREP	59
9.2	Date initialization and default input data file for all schemes	60
9.3	Sea scheme "SEAFLEX"	61

9.4	Lake scheme "WATFLX"	62
9.5	Lake scheme "FLAKE"	63
9.6	Vegetation scheme "ISBA "	64
9.7	Town scheme "TEB "	65
10	How to run the externalized surface physical schemes	67
10.1	"SURF_ATM" general options available over all tiles	67
10.2	"SEAFLX" sea scheme options	68
10.3	"FLAKE" lake scheme options	69
10.4	"ISBA " vegetation scheme options	70
11	How to run the externalized surface chemical schemes	71
11.1	Chemical settings control	71
11.2	Chemical anthropogenic emissions	71
11.3	Chemical deposition over ocean	71
11.4	Chemical deposition over lakes	71
11.5	Chemical deposition over towns	72
11.6	Chemical deposition and biogenic emissions over vegetation	72
12	Externalized surface diagnostics	73
12.1	Diagnostics relative to the general surface monitor	73
12.2	Diagnostics relative to the general surface monitor and to each surface scheme . . .	74
12.3	Diagnostics relative to the ISBA vegetation scheme	75
12.4	Diagnostics relative to the TEB town scheme	75
13	Externalized surface model output fields	77
13.1	Prognostic model output fields	77
13.1.1	ISBA	77
13.1.2	SEAFLUX	79
13.1.3	TEB	80
13.1.4	WATFLUX	80
13.1.5	FLAKE	80
A	Example of namelist	81
A.1	surface parameters defined from ecoclimap	81
A.2	user defined surface parameters	83
A.2.1	Uniform values prescribed : 1d example without patches	83
A.2.2	Uniform values prescribed : 1d example with patches	86
A.2.3	Surface parameters read from external files	94

Première partie

How to install the software

Chapitre 1

Export off-line version of SURFEX

Instructions to install surfex on a linux-PC and to run a 1d example.

0. select a directory where installation has to be done: for example \$HOME or \$HOME/MYDIR, where MYDIR is an existing directory (if not, it has to be created by the user). From now on, it is supposed that the user has defined a MYDIR directory.
1. download EXPORT_v2.tgz from surfex web site and move it into \${HOME}/MYDIR/EXPORT_v2.tgz
2. extract files from archive: tar zxvf EXPORT_v2.tgz
(or gunzip EXPORT_v2.tgz and then tar xvf EXPORT_v2.tar)
at this stage directory EXPORT_v2 is created in MYDIR and contains all software peaces.
3. initialize environment variables needed for surfex:
 - 3.1 set main environment variable:

```
export SURFEX_EXPORT="$HOME/MYDIR/EXPORT_v2"
```
 - 3.2 then run a configuration script included in the archive:

```
. $SURFEX_EXPORT/conf/profile_surfex
```
4. run install_surfex:

```
install_surfex : (i)  realizes the compilation of surfex sources  
                 (ii) prepares executable files for pgd, prep, offline and sxpost applications
```
5. then run install_experiment:

```
install_experiment hapex
```



```
install_experiment : (i) realizes the compilation of prep_input_experiment  
                    (ii) makes useful links to run pgd, prep, offline and sxpost
```
6. once the installation is done, go to \$SURFEX_EXPERIMENT/hapex directory and then to rundir directory and launch successively:
 - (i) pgd.exe
 - (ii) prep.exe
 - (iii) offline.exe

7. to view output 1d, you can use vespa tool. Type vespa to get available fields and then vespa FIELDNAME to plot FIELDNAME

8. How to rerun your own experiment:

8.1 modify \$SURFEX_EXPERIMENT/hapex/rundir/OPTIONS.nam to define new characteristics of PGD and PREP and run pgd.exe and prep.exe (be carefull that diagnostics must be off for PREP application for surfex versions before v3.0)

8.2 modify \$SURFEX_EXPERIMENT/hapex/netcdf/my_forcing.f90 to define your own forcing and parameters of the experiment. Compile the program with sc_input_experiment and run it with prep_input_experiment. Netcdf files (PARAMS.nc and FORCING.nc will be created and need to be placed into \$SURFEX_EXPERIMENT/hapex/rundir directory.

8.3 go to \$SURFEX_EXPERIMENT/hapex/rundir to rerun offline.exe

COMPILING MY OWN SOURCES

```
#
# If you want to compile your "own" sources the preferable
# way is to edit the file "Makefile.SURFEX.mk" in $SURFEX_EXPORT/src.
#
# in the rubric "Source MYSRC" for examples , if your
# sources are in the directory "MY_OWN_SRC" put
```

```
#DIR_MYSRC += MYSRC
DIR_MYSRC += MY_OWN_SRC
```

```
# For the moment , by default your sources will
# be integrated in the original version of the
# already compile librairies with the same
# name ( Not yet BIBUSER management , coming soon ... )
#
# If you want to compile an entire new librarie
# with a new suffix , for examples "MY_MODIF"
# uncomment the lines "ARCH_XYZ.... ",
```

```
VER_MYSRC=MY_MODIF
ARCH_XYZ      := $(ARCH_XYZ)-$(VER_MYSRC)
```

```
# And the suffix "MY_MODIF" will be added to the name of
# the directory of "*.o" and also to the name of the
# libraries ...
```

```
# Once you have edited the "Makefile.SURFEX.mk" file, do the compilation
# using the command:
```

```
make
```

8.4 if you want to modify the source code, go to \$SURFEX_EXPORT/src and do your modifications. Then rebuild executable files by rerunning makefile files. The new executable will be automatically be taken into account in \$SURFEX_EXPERIMENT/hapex/rundir directory.

```
# copy the executables using
#
```

```
ln -s $EXECDIR/pgd.exe      $SURFEX_EXPERIMENT/hapex/rundir/.
ln -s $EXECDIR/prep.exe     $SURFEX_EXPERIMENT/hapex/rundir/.
ln -s $EXECDIR/offline.exe  $SURFEX_EXPERIMENT/hapex/rundir/.
```


Deuxième partie

Off-line guide

Chapitre 2

Using the externalized surface in off-line mode

The use of the externalized surface software in off-line mode requires the preparation of several types of file, especially the input data necessary for the run and the definition of the options specified in the unique namelist . The following example explains how to prepare the different files and how to select the run options.

2.1 Case study

The example is based on the situation of the 25th of October 2004 at 6utc and covers a temporal period of 24 hours. During this day, an unstable weather is observed in France, especially in the Southern part.

2.2 The input files

OPTIONS.nam is the unique namelist used in the off-line mode. It's valid for the PGD, PREP and RUN facilities but for PREP it's necessary to switch off all diagnostics (this is the case by default in surfex). **PARAMS.nc** contains information related to the integration as well as information that is not written in other input files. **FORCING.nc** contains the atmospheric forcing. **PGD.txt** is the PGD file, result of PGD application which prepares physiographical fields. **PREP.txt** is the PREP file containing physiographical information taken from PGD.txt and initial values for prognostic variables of the model extracted from any atmospheric model.

Since ascii and netcdf input files are prepared independantly, one has to take care of the consistency between the number of grid points specified in OPTIONS.nam and the one used in PARAMS.nc, as well as the grid definition used for the forcing data that has to fit the one define by the namelist options.

Chapitre 3

Namelist options

the different namelist options are discussed in the externalized surface user's guide. Here are the main option used for this specific example

3.1 namelist for PGD

- **NAM_PGD_GRID**

CGRID describes the grid type used for the application. In this case it's set to 'LONLAT REG' which means that the working grid will be a regular longitude latitude grid.

- **NAM_LONLAT_REG**

give full definition of the grid by specifying the minimum and maximum values of longitudes and latitudes as well as the number of grid points in each direction.

Variable name	Value	Description
XLONMIN	1.3	
XLONMAX	1.7	
XLATMIN	43.4	
XLATMAX	43.7	
NLON	40	
NLAT	32	

- **NAM_COVER**

YCOVER is the name of the file which contains the ECOCLIMAP database for the definition of the cover present in the working area. **YFILETYPE** is the type of file.

- **NAM_ZS**

YZS is the name of the file which contains the GTOPO30 mnt at 30" resolution (nearly 1km). **YFILETYPE** is the type of file.

- **NAM_ISBA**

YCLAY is the name of the file which contains the clay percentage (for the time being, its taken from FAO at 10 km resolution) **YSAND** is the name of the file which contains the sand percentage (for the time being, its taken from FAO at 10 km resolution)

- **XUNIF_RUNOFFB = 0.5**

sets a uniform value for the slope of the runoff curve

- **CISBA = '3-L'**

corresponds to the standard 3 layers force restore isba scheme

- **CPHOTO = 'NON'**

no specific photosynthesis used in this experiment

- **NPATCH = 1**

single patch for vegetation

- **NGROUND_LAYER = 3**

number of soil layers

3.2 namelist for PREP

- **NAM_PREP_SURF_ATM**
CFILE and **CFILETYPE** are respectively the name and the type of the atmospheric file where fields are read at the level of full surface.
- **NAM_PREP_TEB**
CFILE_TEB and **CTYPE** are respectively the name and the type of the atmospheric file where fields are read for TEB (Town Energy Balance) scheme.
- **NAM_PREP_SEAFLUX**
CFILE_SEAFLX and **CTYPE** are respectively the name and the type of the atmospheric file where fields are read for SEAFLUX scheme.
- **NAM_PREP_WATFLUX**
CFILE_WATFLX and **CTYPE** are respectively the name and the type of the atmospheric file where fields are read for WATERFLUX scheme.
- **NAM_PREP_ISBA**
CFILE_ISBA and **CTYPE** are respectively the name and the type of the atmospheric file where fields are read for ISBA (Interaction between Soil, Biosphere and Atmosphere) scheme.
- **NAM_PREP_ISBA_SNOW**
CSNOW defines the snow scheme used in ISBA.

3.3 namelist for RUN

- **NAM_IO_OFFLINE**
 - **YPROGRAM = 'ASCII'** means that IO will be treated in ascii format and so, PGD and PREP facilities will be used to prepare physiography and surface initial state. Ascii file PREP.txt is expected to initialize physiography and initial values of prognostic fields. The other possibility is **YPROGRAM = 'OFFLIN'**, in that case output of simulations will be written in netcdf files.
 - **YCOUPLING** allows to choose between explicit 'E' and implicit 'I' coupling (It doesn't matter in an off-line simulation, since there's no effective coupling)
 - **YWRPRGM** ascii output format if **YWRPRGM='TEXTE'**, netcdf format if **YWRPRGM='OFFLIN'** (default). Usefull for 1d experiments.
 - **LPRINT** shows the time evolution of your simulation
 - **LRESTART** is used to save the surface state at the end of the simulation. The file that is produced is named SURFOUT.txt (like for PGD) and has to be renamed into PREP.txt to start a new simulation with the saved conditions.
 - **LINQUIRE** enables the inquiry of surface variables from the atmosphere.

Chapitre 4

Data input files

Two different formats are used as input for the externalized surface off-line software. Both are portable : the first format is netcdf and the second is the ascii one. Netcdf format has been chosen because of several participations of PILPS intercomparison projects that requires such format, due to its portability. It follows the Alma concept (proposed by Polcher in 1998).

4.1 netcdf format files

4.1.1 FORCING.nc

For this experiment, atmospheric forcing is extracted from French database named BDAP (Base de Données Analysées et Prévues). Data come from the analysis of surface parameters performed by Safran analysis system devoted to hydrological applications. A constant value in space is applied for each gridbox.

list of parameters

Variable name	Dimensions	Unit	Description
NB_TIMESTP		s	number of time steps
YEAR		s	current year
MONTH			current month
DAY			current day
FORC_TIME_STEP		s	forcing time step
TIME		s	current time
TSUN	DIM_FULL	s	current sun time
ZS	DIM_FULL	m	surface orography
LON	DIM_FULL	deg	longitudes
LAT	DIM_FULL	deg	latitudes
TA	DIM_FULL , NB_TIMESTP	K	air temperature
QA	DIM_FULL , NB_TIMESTP	Kg/Kg	air specific humidity
WIND	DIM_FULL , NB_TIMESTP	m/s	wind speed
DIR_SW	DIM_FULL , NB_TIMESTP	W/m2	downward direct shortwave radiation
SCA_SW	DIM_FULL , NB_TIMESTP	W/m2	downward diffuse shortwave radiation
LW	DIM_FULL , NB_TIMESTP	W/m2	downward longwave radiation
PS	DIM_FULL , NB_TIMESTP	Pa	surface pressure
RAIN	DIM_FULL , NB_TIMESTP	Kg/m2/s	rainfall rate
SNOW	DIM_FULL , NB_TIMESTP	Kg/m2/s	snowfall rate
CO2	DIM_FULL , NB_TIMESTP		CO2 concentration
DIR	DIM_FULL , NB_TIMESTP	deg	wind direction

Dimensions **DIM_FULL** and **NB_TIMESTP** represent respectively the total number of gridboxes in the area of interest and the number of atmospheric time steps.

The following pictures show the time evolution of forcing quantities for the integration period, over the region of interest. Each curve correspond to a grid point.




FIG. 4.1 – *Time evolution of temperature over the working area*

4.1.2 PARAMS.nc

Variable name	Dimensions	Value	Unit	Description
ISBA_TIME_STEP		300	s	isba time step
OUTPUT_TIME_STEP		3600	s	time step for writing output
DIM_FULL		1280		number of grid boxes
GROUND_LAYER		3		number of layers in the soil
SNOW_LAYER		3		number of snow layers
PATCH_NUMBER		1		number of patch
DTCUR-TDATE		2004, 10, 25		starting date of the run
DTCUR-TIME		21600	s	starting time of the run
ZREF	DIM_FULL	2	m	reference height for T and q
UREF	DIM_FULL	10	m	reference height for wind

The two netcdf files are prepared independantly from PGD and PREP applications, with external programs named `prep_input_parameters.f90` and `my_forcing.f90`. Parameters `DIM_FULL`, `GROUND_LAYER`, `SNOW_LAYER`, `PATCH_NUMBER`, `DTCUR-TDATE`, `DTCUR-TIME` have to be set up carefully according to the one defined during PGD and PREP applications, anyhow consistency tests are performed during surfex setup.

4.2 ascii format files

This format is used in the off-line applications because it's the simplest format that may replace more complex (in terms of file handling) like lfi format used in the meso-NH framework or FA used in the Arome framework.




FIG. 4.2 – *Time evolution of specific humidity over the working area*




FIG. 4.3 – *Time evolution of longwave incoming radiation over the working area*




FIG. 4.4 – *Time evolution of shortwave incoming radiation over the working area*

4.2.1 PGD.txt

This file contains the information related to physiography and orography essentially. The file is splitted into several parts. The first one corresponds to the gridbox as seen as a single pixel where quantities are aggregated. The corresponding field names are prefixed with **FULL**. This gridbox may be separated into four tiles respectively associated to nature, town, sea/ocean and lake). The corresponding field names are respectively prefixed with **NATURE**, **TOWN**, **SEA** and **WATER**.

The physiographic parameters written out into PGD.txt file are mainly the fraction of land covers contained in each gridbox. These fractions are computed from ECOCLIMAP database.

```
&FULL DIM_FULL
(-)
1280
&FULL DIM_SEA
(-)
0
&FULL DIM_NATURE
(-)
1279
&FULL DIM_WATER
(-)
1
&FULL DIM_TOWN
(-)
240
&FULL ECOCLIMAP
(-)
T
&NATURE ISBA
ISBA
3-L
&NATURE PHOTO
PHOTO
NON
&NATURE GROUND_LAYER
GROUND_LAYER
3
&NATURE PATCH_NUMBER
PATCH_NUMBER
1
```

Value of **DIM_FULL** indicates that this is a 2D exercise, and the gridbox contains a non-zero fraction of nature, of water (lake) and of town, but there's no fraction of sea because the number of points with a non-zero fraction of sea **DIM_SEA** is zero. 1279 gridboxes contain vegetation and 240 gridboxes contain a fraction of town, onmly one gridbow contains a fraction ok inland water(lake). Surface scheme to treat vegetation is Isba 3-L which means that soil is represented with 3 layers. The number of patches is 1, it indicates that the vegetation is not splitted into patches like it could (this should be the case for the A-gs option of Isba that treats explicitly photosynthesis).




FIG. 4.5 – *Orography field taken from PGD.txt file*

4.2.2 PREP.txt

This file contains the information related to physiography and orography read from PGD.txt file and rewritten, as well as initial values of prognostic variables of the different schemes contained in SURFEX.

```
&NATURE TG1
X_Y_TG1 (K)
  2.8576409563069382E+02  2.8548082006251650E+02  2.8540527530138650E+02  2.8546873415878122E+02 ...
```

For example here are shown the first initial values of skin surface temperature over natural area for Isba surface scheme as they are written in PREP.txt file. This field is represented on figure 4.6.




FIG. 4.6 – *Initial surface temperature for vegetation taken from PREP.txt file*

An other example shows the roof surface temperature over the working area (Toulouse city is located roughly at $x=15$, $y=22$)

4.2.3 Extracting 2d fields

It's possible to extract 2d fields from PGD.txt (covers, orography, etc.) and from PREP.txt (initial prognostic variables like soil temperature profile, soil water content profile, etc.). For that purpose you need to run SXPOST tool (located at the same place as PGD, PREP and OFFLINE : \$SURFEX_EXPORT/src/exe/). Input files for SXPOST are PGD.txt or PREP.txt if exists, and a namelist




FIG. 4.7 – Initial surface temperature for the roofs taken from *PREP.txt* file

containing the number of fields to be extracted, the name and a flag indicating if the variable depends on patches or not. The name of a given field is the name written in PGD.txt or PREP.txt file where characters " &" have been removed. For example, to extract orography, the name of the field is ZS in SURFEX, the mask over which it's defined can be FULL (total gridbox). To extract surface temperature over vegetation, the name will be TG1 and the mask NATURE. (grep "&" PGD.txt returns all variables of this file). To distinguish variables defined or not over patches, a flag is used : if the variable is patch dependant, the flag must be set to '+', in the contrary, it should be '-'. For example orography (ZS) doesn't depend on patches, but surface temperature (TG1) does. If the simulation uses patches and the flag is '-' then only the first patch will be treated (bare ground).

The namelist SXPOST.nam looks like :

```
2
- FULL   ZS
+ NATURE TG1
```

Running SXPOST will return a file per variable, which will contain the longitude, the latitude and the value of the field for each gridbox over which the field is defined (For example, TG1 which is known only over nature won't have a value for each gridbox of the domain).

4.2.4 I/O diagram




FIG. 4.8 – Surfex diagram showing the input/output files produced by the different tools (the list of produced variables in case **YWRPRGM='TEXTE'** is obviously not exhaustive)

Chapitre 5

Architecture of the off-line driver

5.1 Initialization

At first, the driver makes some initializations. For example, physics constants, geoid description, etc. are initialized through **ini_csts** subroutine. Then information is read from input files **PARAMS.nc** and **PREP.txt** in subroutine **ol_read_atm_conf**, it concerns especially the run configuration : time step for surface, time step for the atmosphere, dimension of the grid, date, etc. It reads also latitude and longitude in **FORCING.nc** file and realizes some check consistency between input information (netcdf files and **PREP.txt** file).

After that, the atmospheric variables are initialized : air temperature, specific humidity, wind speed and direction, shortwave (direct and scattered) and longwave incoming radiation, liquid and solid precipitation, CO2 concentration and surface pressure.

Then the main part of the initialization **init_surf_atm** is called : a description is given in "Overview of SURFEX setup (Le Moigne)".

5.2 Temporal loop

```
DO JFORC_STEP=1,INB_STEP_ATM
  !
  ! read Forcing
  !
  CALL OL_READ_ATM(YPROGRAM, JFORC_STEP,
                  ZTA,ZQA,ZWIND,ZDIR_SW,ZSCA_SW,ZLW,ZSNOW,ZRAIN,ZPS,&
                  ZCO2,ZDIR
                  )

DO JSURF_STEP=1,INB_ATM
  !
  ! time interpolation of the forcing
  !
  CALL SUNPOS(IYEAR, IMONTH, IDAY, ZTIME, ZLON, ZLAT, XTSUN, XZENITH, XAZIM)

  CALL OL_TIME_INTERP_ATM(JSURF_STEP,INB_ATM,
                        ZTA,ZQA,ZWIND,ZDIR_SW,ZSCA_SW,ZLW,ZSNOW,ZRAIN,ZPS,&
                        ZCO2,ZDIR
                        )
```

There are 2 main loops : the external (JFORC_STEP) concerns the atmosphere and the internal deals with the surface (JSURF_STEP). For each atmospheric step, the forcing variables are read from **FORCING.nc**. In fact, 2 time steps are read because, during the surface loop, forcing are linearly interpolated in time (**ol_time_interp_atm**) because the surface time step is shorter than the atmosphere one. Once all forcing are known for a given surface step, physical computation may begin.

5.2.1 coupling surface with the atmosphere

```
CALL COUPLING_SURF_ATM_n(YPROGRAM, YCOUPLING,           &
  ZTSTEP_ISBA, IYEAR, IMONTH, IDAY, ZTIME, INI, ISCAL, IBANDS, &
  XTSUN, XZENITH, XAZIM,                                &
  XZREF, XUREF, XZS, XU, XV, XQA, XTA, XROA, XSV, XCO2, CSV, &
  XRAIN, XSNOW, XLW, XDIR_SW, XSCA_SW, XSW_BANDS, XPS, XPA, &
  XSFTQ, XSFTH, XSFTS, XSFCO2, XSFU, XSFV,             &
  XTSRAD, XDIR_ALB, XSCA_ALB, XEMIS,                   &
  XPEW_A_COEF, XPEW_B_COEF,                             &
  XPET_A_COEF, XPEQ_A_COEF, XPET_B_COEF, XPEQ_B_COEF, YTEST )
```

coupling_surf_atm is the subroutine that realizes the interface between the surface and the atmosphere. It receives the atmospheric forcing, some configuration parameters (time step, dimensions), and returns the fluxes computed at surface, aggregated for each gridbox.

5.2.2 writing out prognostic variables

```
CALL WRITE_SURF_ATM_n('OFFLIN', 'ALL')
```

This subroutine writes out the prognostic variables of the different schemes. In off-line mode, they will be written out into netcdf files named ISBA_PROGNOSTIC.OUT.nc, TEB_PROGNOSTIC.OUT.nc, SEAFLUX_PROGNOSTIC.OUT.nc, WATFLUX_PROGNOSTIC.OUT.nc. Some diagnostics may also be written (depending on activated switch in namelist OPTIONS.nam)

5.2.3 compute diagnostics

```
CALL DIAG_SURF_ATM_n('OFFLIN')
```

This subroutine computes diagnostics that have not been computed during the the time step evolution (coupling_surf_atm computes automatically some diagnostics : 2 meters parameters, 10 meter wind, and all fluxes that will be return to the atmosphere when running in a coupled way). These computation is decided through namelist OPTIONS.nam (see User's guide for full description).

5.2.4 writing out diagnostics variables

```
CALL WRITE_DIAG_SURF_ATM_n('OFFLIN', 'ALL')
```

Once these diagnostics have been computed, they are stored in modules and will be effectively written to files when calling write_diag_surf_atmn subroutine. The possible diagnostic files are ISBA_DIAGNOSTICS.OUT.nc, TEB_DIAGNOSTICS.OUT.nc, SEAFLUX_DAIGNOSTICS.OUT.nc, WATFLUX_DIAGNOSTICS.OUT.nc and SURF_ATM_DIAGNOSTICS.OUT.nc. This last file contains aggregated fields over the whole gridbox.

5.3 writing out restart file

```
!
!*   3.   write restart file
!       -----
!
IF ( LRESTART ) CALL WRITE_SURF_ATM_n('ASCII ', 'ALL')
```

This file will be name SURFOUT.txt and will content the surface state at the end of the off-line run. It ishould replace the PREP.txt file in case of restart of the model.

Chapitre 6

Some output of off-line simulation

6.1 Examples of prognostic variables output




FIG. 6.1 – *Initial surface temperature field*




FIG. 6.2 – *Root layer water content after 12 hours of integration*




FIG. 6.3 – *Interception (by vegetation) water content after 12 hours of integration*

6.2 list of available variables

This list has been made by using vespa which is a tool that has been used to realize the plots of this document. This is not the complete possible list since only the files that are present in netcdf output files are listed.

```
*****
***** V E S P A *****
*****
***          ***
*** V-aluable ***
*** E-externalized ***
*** S-urface ***
*** P-lot ***
*** A-pplication ***
***          ***
*****
*****

ISBA PROGNOSTIC VARIABLES
=====

TG1: "Soil_temp_layer_1" ;
WG1: "Soil_liquid_layer_1" ;
WG11: "Soil_ice1" ;
TG2: "Soil_temp_layer_2" ;
WG2: "Soil_liquid_layer_2" ;
WG12: "Soil_ice2" ;
TG3: "Soil_temp_layer_3" ;
WG3: "Soil_liquid_layer_3" ;
WG13: "Soil_ice3" ;
WR: "Interception_reservoir" ;
RESA: "Aerodynamic_resistance" ;
WSNOW_VEG1: "Snow_Water_Equivalent_layer_1" ;
RSNOW_VEG1: "Snow_density_layer_1" ;
TSNOW_VEG1: "Snow_temperature_layer1" ;
HSNOW_VEG1: "Snow_heat_layer1" ;
WSNOW_VEG2: "Snow_Water_Equivalent_layer_2" ;
RSNOW_VEG2: "Snow_density_layer_2" ;
TSNOW_VEG2: "Snow_temperature_layer2" ;
HSNOW_VEG2: "Snow_heat_layer2" ;
WSNOW_VEG3: "Snow_Water_Equivalent_layer_3" ;
RSNOW_VEG3: "Snow_density_layer_3" ;
TSNOW_VEG3: "Snow_temperature_layer3" ;
HSNOW_VEG3: "Snow_heat_layer3" ;
ASNOW_VEG: "Snow_albedo" ;

ISBA DIAGNOSTIC VARIABLES
=====

RI_ISBA: "Averaged_Richardson_Number" ;
RN_ISBA: "Averaged_Net_Radiation" ;
H_ISBA: "Averaged_Sensible_Heat_Flux" ;
LE_ISBA: "Averaged_Latent_Heat_Flux" ;
GFLUX_ISBA: "Averaged_Ground_Heat_Flux" ;
LEG: "Ground_Evaporation_Heat_Flux" ;
LEGI: "Soil_Ice_Sublimation" ;
LEV: "Vegetation_Evaporation_Heat_Flux" ;
LES: "Snow_Evaporation_Heat_Flux" ;
LER: "Canopy_Water_Interception_Evaporation" ;
LETR: "Vegetation_Evapotranspiration" ;
EVAP: "Evapotranspiration" ;
DRAIN: "Soil_Drainage_Flux" ;
RUNOFF: "Supersaturation_Runoff" ;
LEG_ISBA: "Averaged_Ground_Evaporation_Heat_Flux" ;
LEGI_ISBA: "Averaged_Soil_Ice_Sublimation" ;
LEV_ISBA: "Averaged_Vegetation_Evaporation_Heat_Flux" ;
LES_ISBA: "Averaged_Snow_Evaporation_Heat_Flux" ;
LER_ISBA: "Averaged_Canopy_Water_Interception_Evaporation" ;
LETR_ISBA: "Averaged_Vegetation_Evapotranspiration" ;
EVAP_ISBA: "Averaged_Evapotranspiration" ;
DRAIN_ISBA: "Averaged_Soil_Drainage_Flux" ;
RUNOFF_ISBA: "Averaged_Supersaturation_Runoff" ;
CH_ISBA: "Averaged_thermal_diffusion_coefficient" ;
HV_ISBA: "Halstead_coefficient" ;
ZOREL: "Output_ZOREL" ;
VEGTYPE_PATCH_1: "fraction_of_vegetation_type_1" ;
VEGTYPE_PATCH_2: "fraction_of_vegetation_type_2" ;
VEGTYPE_PATCH_3: "fraction_of_vegetation_type_3" ;
VEGTYPE_PATCH_4: "fraction_of_vegetation_type_4" ;
VEGTYPE_PATCH_5: "fraction_of_vegetation_type_5" ;
VEGTYPE_PATCH_6: "fraction_of_vegetation_type_6" ;
VEGTYPE_PATCH_7: "fraction_of_vegetation_type_7" ;
VEGTYPE_PATCH_8: "fraction_of_vegetation_type_8" ;
VEGTYPE_PATCH_9: "fraction_of_vegetation_type_9" ;
VEGTYPE_PATCH_10: "fraction_of_vegetation_type_10" ;
VEGTYPE_PATCH_11: "fraction_of_vegetation_type_11" ;
VEGTYPE_PATCH_12: "fraction_of_vegetation_type_12" ;

ISBA PHYSIOGRAPHIC VARIABLES
=====

VEG: "Output_vegetation_fraction" ;
ZO_ISBA: "Output_ZO_ISBA" ;
LAI: "Output_LAI_ISBA" ;
ALBNIR_SOIL: "Output_ALBNIR_SOIL" ;
ALBVIS_SOIL: "Output_ALBVIS_SOIL" ;
```

TEB PROGNOSTIC VARIABLES

=====

```
T_ROOF1: "Roof_Temperature_Layer_1" ;
T_ROOF2: "Roof_Temperature_Layer_2" ;
T_ROOF3: "Roof_Temperature_Layer_3" ;
T_ROAD1: "Road_Temperature_Layer_1" ;
T_ROAD2: "Road_Temperature_Layer_2" ;
T_ROAD3: "Road_Temperature_Layer_3" ;
T_WALL1: "Wall_Temperature_Layer_1" ;
T_WALL2: "Wall_Temperature_Layer_2" ;
T_WALL3: "Wall_Temperature_Layer_3" ;
TI_BLD: "Internal_Building_Temperature" ;
TI_ROAD: "Deep_Road_Temperature" ;
WS_ROOF1: "Roof_Water_Content_Layer_1" ;
WS_ROOF2: "Roof_Water_Content_Layer_2" ;
WS_ROOF3: "Roof_Water_Content_Layer_3" ;
WS_ROAD1: "Road_Water_Content_Layer_1" ;
WS_ROAD2: "Road_Water_Content_Layer_2" ;
WS_ROAD3: "Road_Water_Content_Layer_3" ;
T_CANYON: "Canyon_Air_Temperature" ;
Q_CANYON: "Canyon_Air_Humidity" ;
```

TEB DIAGNOSTIC VARIABLES

=====

```
RI_TEB: "Averaged_Richardson_Number" ;
CD_TEB: "Averaged_Drag_Momentum_Coef" ;
CDN_TEB: "Averaged_Neutral_Drag_Coef" ;
CH_TEB: "Averaged_Drag_Thermal_Coef" ;
RESA_TEB: "Averaged_Aerodyn_Resistance" ;
RN_TEB: "Averaged_Net_Radiation" ;
H_TEB: "Averaged_Sensible_Heat_Flux" ;
LE_TEB: "Averaged_Latent_Heat_Flux" ;
GFLUX_TEB: "Averaged_Ground_Heat_Flux" ;
```

WATER PROGNOSTIC VARIABLES

=====

```
TS_WATER: "Averaged_Water_Temperature" ;
ZO_WATER: "Roughness length" ;
```

WATER DIAGNOSTIC VARIABLES

=====

```
RI_WAT: "Averaged_Richardson_Number" ;
CD_WAT: "Averaged_Drag_Momentum_Coef" ;
CDN_WAT: "Averaged_Neutral_Drag_Coef" ;
CH_WAT: "Averaged_Drag_Thermal_Coef" ;
RESA_WAT: "Averaged_Aerodyn_Resistance" ;
RN_WAT: "Averaged_Net_Radiation" ;
H_WAT: "Averaged_Sensible_Heat_Flux" ;
LE_WAT: "Averaged_Latent_Heat_Flux" ;
GFLUX_WAT: "Averaged_Ground_Heat_Flux" ;
```

SURF_ATM DIAGNOSTICS VARIABLES

=====

```
RI: "Averaged_Richardson_Number" ;
RN: "Averaged_Net_Radiation" ;
H: "Averaged_Sensible_Heat_Flux" ;
LE: "Averaged_Latent_Heat_Flux" ;
GFLUX: "Averaged_Ground_Heat_Flux" ;
```

SURF_ATM FRACTIONS

=====

```
FRAC_SEA: "Fraction_of_sea" ;
FRAC_WATER: "Fraction_of_water" ;
FRAC_TOWN: "Fraction_of_town" ;
FRAC_NATURE: "Fraction_of_nature" ;
```

FORCING FIELDS

=====

```
TA: "air temperature" ;
QA: "air specific humidity" ;
WIND: "wind speed" ;
DIR_SW: "downward direct shortwave radiation" ;
SCA_SW: "downward diffuse shortwave radiation" ;
LW: "downward longwave radiation" ;
PS: "surface pressure" ;
RAIN: "rainfall rate" ;
SNOW: "snowfall rate" ;
CO2: "CO2 concentration" ;
DIR_SW: "downward direct shortwave radiation" ;
DIR: "wind direction" ;
```


Troisième partie

Users guide

Chapitre 7

Overview of the externalized surface sequence

The externalized surface facilities do not contain only the program to run the physical surface schemes, but also those producing the initial surface fields (before the run) and the diagnostics (during or after the run). All these facilities are listed, below, and they separate in 4 main parts :

7.1 The sequence

1. **PGD** (routine `pgd_surf_atm.n.f90`) : this program computes the physiographic data file (called PGD file below). At this step, you perform 3 main tasks :
 - (a) You choose the surface schemes you will use.
 - (b) You choose and define the grid for the surface
 - (c) The physiographic fields are defined on this grid.

Therefore, the PGD file contains the spatial characteristics of the surface and all the physiographic data necessary to run the interactive surface schemes for vegetation and town.

2. **PREP** (routine `prep_surf_atm.n.f90`) : this program performs the initialization of the surface scheme prognostic variables, as temperatures profiles, water and ice soil contents, interception reservoirs, snow reservoirs.
3. **run of the schemes** (routine `coupling_surf_atm.n.f90`) : this performs the physical evolution of the surface schemes. It is necessary that this part, contrary to the 2 previous ones, is to be coupled within an atmospheric forcing (provided either in off-line mode or via a coupling with an atmospheric model).
4. **DIAG** (routine `diag_surf_atm.n.f90`) : this computes diagnostics linked to the surface (e.g. surface energy balance terms, variables at 2m of height, etc...). It can be used either during the run (adding these diagnostics in the output file(s) of the run), or independantly from the run, for a given surface state (still, an instantaneous atmospheric forcing is necessary for this evaluation).

In addition, in order to read or write the prognostic variables or the diagnostics variables, respectively, in the surface files, the following subroutines are used : `init_surf_atm.n.f90`, `write_surf_atm.n.f90` and `write_diag_surf_atm.n.f90`.

7.2 The atmospheric models using the externalized surface

The externalized surface can presently be used in :

1. in offline mode
2. MESONH
3. AROME

For each model, additionnal possibilities of the surface, especially the ability to read and write in files with particular formats, are added :

7.2.1 in offline mode

In this case, 2 types of files can be used :

- *ASCII files*, not efficient in term of storage, but completely portable.
- *netcdf files*, that can be used by the program code "OFFLIN".

currently, **PGD** and **PREP** steps are done using ASCII files, while the run produces time series of each variable (prognostic or diagnostic) in netcdf files and the output instant of the run again in an ASCII file.

The nameists are all included in the namelist file named `OPTIONS.nam`

7.2.2 in MESONH

In this case, MESONH FM files are used. The parallelization of the surface fields is done during the reading or writing of the fields by the `FMREAD` and `FMWRIT` routines.

Initialization of surface fields integrated in MESONH programs

In MESONH, there are usually 2 ways to produce initial files, depending if you want to use real or ideal atmospheric conditions. However, from the surface point of view, there is no difference between these 2 main possibilities of fields (real -e.g. from operationnal surface scheme in an operationnal model- or ideal -e.g. uniform-), whatever the treatment done for the atmospheric fields. This is allowed because the same externalized routines corresponding to **PGD** and **PREP** are used :

In the case of realistic atmospheric fields, the MESONH programs calling the surface are :

1. **PREP_PG**D : it uses the **PGD** facility of the surface
2. **PREP_NEST_PG**D : surface fields are only read and rewritten, except the orography that is modified (the modification of the orography itself is considered as an atmospheric model routine, as orography is also a field of the atmospheric model).
3. **PREP_REAL**CASE : it uses the **PREP** facility of the surface, that can produce either ideal or realistic surface fields.

4. SPAWNING : it does not produce surface fields any more. The surface fields will be recreated during the PREP_REAL_CASE step following the SPAWNING.

In the case of ideal atmospheric fields, the MESONH program calling the surface is :

1. PREP_IDEAL_CASE : it uses both the **PGD** and **PREP** facilities of the surface. Ideal or realistic (the latter only in conformal projection) physiographic fields can be either produced or read from a file. Then the prognostic surface variables, either ideal or realistic, can be computed by **PREP**.

If you use MESONH atmospheric model, the input and output surface files are the same as the atmospheric ones, so there is no need to specify via surface namelists any information about the input or output file names.

Namelist NAM_PGDFILE

Note however that, in PREP_PGDF (just before the call to the surface physiographic computation in **PGD**, for which the namelists are described in the next chapter), there is a namelist to define the output physiographic file :

Fortran name	Fortran type
CPGDFILE	string of 28 characters

MESONH run and diagnostics

Then, the MESONH run can be done. During this one, the diagnostics can be, or not, be computed.

In DIAG, the surface diagnostics can also be recomputed.

7.2.3 in AROME

In this case, MESONH FM files are also used, for the surface only. The parallelization of the surface fields is done during the reading or writing of the fields by parallelization routines of ALADIN atmospheric model.

Chapitre 8

The physiographic fields

8.1 Overview of physiographic fields computation : PGD

The physiographic fields are averaged or interpolated on the specified grid by the program **PGD**. They are stored in a file, called PGD file, but only with the physiographic 2D fields, the geographic and grid data written in it.

During the **PGD** facility :

1. You choose the surface schemes you will use.
2. You choose and define the grid for the surface.
3. The physiographic fields are defined on this grid.

8.1.1 Choice of the grid

There are 3 possibilities. 2 are always possible, one is available only if the **PGD** routine is integrated into an atmospheric model initialization facility.

1. The grid is chosen via namelists options (see below)
2. The grid is defined as a part of the grid of an already existing surface file, indicated via namelists (see below)
3. The grid is defined as being identical to the one of an atmospheric model, which is given as fortran argument in the coupling of the **PGD** surface facilities (routine **PGD_SURF_ATM**) into an atmospheric model initialization procedures. In this case, **all namelists that are usually used to define the surface grid are ignored**. Note that, in addition to the grid, the orography can also be given from the atmospheric file.

8.1.2 Choice of the physiographic fields

There are 3 main possibilities depending on **LECOCLIMAP** flag.

Namelist **NAM_FRAC**

This namelist defines if **ECOCLIMAP** mechanism based on fractions of covers will be used or not.

Fortran name	Fortran type	values	default value
LECOCLIMAP	Logical		.TRUE.
XUNIF_SEA	real	between 0 and 1	none
CFNAM_SEA	character (LEN=28)		','
CFTYP_SEA	character (LEN=6)	'DIRECT', 'BINLLF', 'BINLLV', 'ASCLLV'	none
XUNIF_WATER	real	between 0 and 1	none
CFNAM_WATER	character (LEN=28)		','
CFTYP_WATER	character (LEN=6)	'DIRECT', 'BINLLF', 'BINLLV', 'ASCLLV'	none
XUNIF_NATURE	real	between 0 and 1	none
CFNAM_NATURE	character (LEN=28)		','
CFTYP_NATURE	character (LEN=6)	'DIRECT', 'BINLLF', 'BINLLV', 'ASCLLV'	none
XUNIF_TOWN	real	between 0 and 1	none
CFNAM_TOWN	character (LEN=28)		','
CFTYP_TOWN	character (LEN=6)	'DIRECT', 'BINLLF', 'BINLLV', 'ASCLLV'	none

- **XUNIF_SEA** : uniform prescribed value of sea fraction. If **XUNIF_SEA** is set, file **CFNAM_SEA** is not used.
- **CFNAM_SEA** : sea fraction data file name. If **XUNIF_SEA** is set, file **CFNAM_SEA** is not used.
- **CFTYP_SEA** : type of sea data file ('DIRECT', 'BINLLF', 'BINLLV', 'ASCLLV')
- **XUNIF_WATER** : uniform prescribed value of water fraction. If **XUNIF_WATER** is set, file **CFNAM_WATER** is not used.
- **CFNAM_WATER** : water fraction data file name. If **XUNIF_WATER** is set, file **CFNAM_WATER** is not used.
- **CFTYP_WATER** : type of water data file ('DIRECT', 'BINLLF', 'BINLLV', 'ASCLLV')
- **XUNIF_NATURE** : uniform prescribed value of nature fraction. If **XUNIF_NATURE** is set, file **CFNAM_NATURE** is not used.
- **CFNAM_NATURE** : nature fraction data file name. If **XUNIF_NATURE** is set, file **CFNAM_NATURE** is not used.
- **CFTYP_NATURE** : type of nature data file ('DIRECT', 'BINLLF', 'BINLLV', 'ASCLLV')
- **XUNIF_TOWN** : uniform prescribed value of town fraction. If **XUNIF_TOWN** is set, file **CFNAM_TOWN** is not used.
- **CFNAM_TOWN** : town fraction data file name. If **XUNIF_TOWN** is set, file **CFNAM_TOWN** is not used.
- **CFTYP_TOWN** : type of town data file ('DIRECT', 'BINLLF', 'BINLLV', 'ASCLLV')

If flag **LECOCLIMAP** is set to .TRUE., there are 2 possibilities :

ideal physiographic fields :

These fields are either uniform (fraction of each 215 ecoclimap ecosystem, orography, or any field needed by the surface schemes). As mentioned above, orography can be, in the case of the coupling with an atmospheric model, imposed as the atmospheric model (non-uniform) field.

realistic physiographic fields :

PGD can use files to build accurate physiographic fields from geographical information. This is possible only if the grid chosen can be linked to geographical coordinates (latitude and longitude), i.e. if the grid type is "LATLONREG " or "CONF PROJ ".

The files that can be used are :

- A file describing the type of cover of the surface. This describes where are located the different cover types (forests, towns, seas, etc...). At the time being, the file provided contains the ecoclimap data (215 land covers) on the world, with a resolution of 30". **PGD** computes the fraction of surface coverage occupied by each type in the grid mesh. From this information, the surface parameter convenient for the surface schemes (such as building fraction, leaf area index, etc..) are deduced, using correspondance arrays : **a parameter has always the same value for a given cover type, anywhere in the world.**
- A file containing the orography : GTOPO30. The resolution of the file is 30" on the world. This allows to compute the model orography, and the following subgrid-scale orographic characteristics :
 - the surface of frontal obstacle (A) over the surface of the grid mesh (S) in each direction ($\sum A_{i+}/S, \sum A_{i-}/S, \sum A_{j+}/S, \sum A_{j-}/S$, used to compute the directional z_{0eff}),
 - the half height of these obstacles ($h_i^+/2, h_i^-/2, h_j^+/2, h_j^-/2$, used to compute the directional z_{0eff}),
 - These 8 parameters are used to compute the total roughness length in the four directions given by the model axis ($z_{0eff_{i+}}, z_{0eff_{i-}}, z_{0eff_{j+}}, z_{0eff_{j-}}$),
 - the Subgrid-Scale Orography (SSO) parameters (standard deviation μ_{z_s} , anisotropy γ_{z_s} , direction of the small main axis θ_{z_s} and slope σ_{z_s}).
- For ISBA scheme, a file with the clay fraction of the (near-surface) soil. The resolution of the file provided is 5' on the world.
- For ISBA scheme, a file with the sand fraction of the (near-surface) soil. The resolution of the file provided is 5' on the world.

If **LECOCLIMAP** flag is set to .FALSE. :

user defined physiographic fields :

ISBA scheme

Over natural areas, all surface parameters for each patch, at a given frequency have to be specified by the user in namelist **NAM_DATA_ISBA**.

parameters depending on the number of vegetation types :

Fortran name	Fortran type	values	default value	description	unit
XUNIF_VEGTYPE	real	between 0 and 1	none	vegetation type	
CFNAM_VEGTYPE	character (LEN=28)		' '	file name	
CFTYP_VEGTYPE	character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none	file type	

parameters depending on the number of patches and time :

Fortran name	Fortran type	values	default value	description	unit
NTIME	integer	12 or 36	36	time dimension	
XUNIF_VEG	real	between 0 and 1	none	vegetation fraction	(-)
CFNAM_VEG	character (LEN=28)		' '	file name	
CFTYP_VEG	character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none	file type	
XUNIF_LAI	real		none	leaf area index	(m^2/m^2)
CFNAM_LAI	character (LEN=28)		' '	file name	
CFTYP_LAI	character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none	file type	
XUNIF_Z0	real		none	roughness length	(m)
CFNAM_Z0	character (LEN=28)		' '	file name	
CFTYP_Z0	character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none	file type	
XUNIF_EMIS	real		none	emissivity	(-)
CFNAM_EMIS	character (LEN=28)		' '	file name	
CFTYP_EMIS	character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none	file type	

parameters depending on the number of patches and soil levels :

Fortran name	Fortran type	values	default value	description	unit
XUNIF_DG	real		none	soil layer thickness	(m)
CFNAM_DG	character (LEN=28)		' '		
CFTYP_DG	character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none		
XUNIF_ROOTFRAC	real		none	root fraction	(-)
CFNAM_ROOTFRAC	character (LEN=28)		' '		
CFTYP_ROOTFRAC	character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none		

parameters depending on number of patches only :

Fortran name	Fortran type	values	default value	description	unit
XUNIF_RSMIN	real		none	minimal stomatal resistance	(s/m)
CFNAM_RSMIN	character (LEN=28)		' '		
CFTYP_RSMIN	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			
XUNIF_GAMMA	real		none	coefficient used in the computation of RSMIN	(-)
CFNAM_GAMMA	character (LEN=28)		' '		
CFTYP_GAMMA	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			
XUNIF_WRMAX_CF	real		none	coefficient for maximum interception water storage capacity	(-)
CFNAM_WRMAX_CF	character (LEN=28)		' '		
CFTYP_WRMAX_CF	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			
XUNIF_RGL	real		none	maximum solar radiation available for photosynthesis	(W/m^2)
CFNAM_RGL	character (LEN=28)		' '		
CFTYP_RGL	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			
XUNIF_CV	real		none	vegetation thermal inertia coefficient	(Km^2/J)
CFNAM_CV	character (LEN=28)		' '		
CFTYP_CV	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			
XUNIF_Z0_O_Z0H	real		none	ratio of surface roughness lengths	(-)
CFNAM_Z0_O_Z0H	character (LEN=28)		' '		
CFTYP_Z0_O_Z0H	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			
XUNIF_ALBNIR_VEG	real		none	vegetation near-infra-red albedo	(-)
CFNAM_ALBNIR_VEG	character (LEN=28)		' '		
CFTYP_ALBNIR_VEG	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			
XUNIF_ALBVIS_VEG	real		none	vegetation visible albedo	(-)
CFNAM_ALBVIS_VEG	character (LEN=28)		' '		
CFTYP_ALBVIS_VEG	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			
XUNIF_ALBUV_VEG	real		none	vegetation UV albedo	(-)
CFNAM_ALBUV_VEG	character (LEN=28)		' '		
CFTYP_ALBUV_VEG	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			
XUNIF_ALBNIR_SOIL	real		none	soil near-infra-red albedo	(-)
CFNAM_ALBNIR_SOIL	character (LEN=28)		' '		
CFTYP_ALBNIR_SOIL	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			
XUNIF_ALBVIS_SOIL	real		none	soil visible albedo	(-)
CFNAM_ALBVIS_SOIL	character (LEN=28)		' '		
CFTYP_ALBVIS_SOIL	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			
XUNIF_ALBUV_SOIL	real		none	soil UV albedo	(-)
CFNAM_ALBUV_SOIL	character (LEN=28)		' '		
CFTYP_ALBUV_SOIL	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			

Fortran name	Fortran type	values	default	description	unit
XUNIF_GMES CFNAM_GMES CFTYP_GMES	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	mesophyll conductance	(m/s^{-1})
XUNIF_BSLAI CFNAM_BSLAI CFTYP_BSLAI	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	ratio d(biomass)/d(lai)	(kg/m^2)
XUNIF_LAIMIN CFNAM_LAIMIN CFTYP_LAIMIN	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	minimum LAI	(m^2/m^2)
XUNIF_SEFOLD CFNAM_SEFOLD CFTYP_SEFOLD	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	e-folding time for senescence	(s)
XUNIF_GC CFNAM_GC CFTYP_GC	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	cuticular conductance	(m/s)
XUNIF_DMAX CFNAM_DMAX CFTYP_DMAX	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	maximum air saturation deficit	(kg/kg)
XUNIF_F2I CFNAM_F2I CFTYP_F2I	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	critical normilized soil water content for stress parameterization	(-)
XUNIF_H_TREE CFNAM_H_TREE CFTYP_H_TREE	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	height of trees	(m)
XUNIF_RE25 CFNAM_RE25 CFTYP_RE25	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	Ecosystem respiration parameter	$(kg/kgms^{-1})$
XUNIF_CE_NITRO CFNAM_CE_NITRO CFTYP_CE_NITRO	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	leaf aera ratio sensivity to [nitrogen]	(m^2/kg)
XUNIF_CF_NITRO CFNAM_CF_NITRO CFTYP_CF_NITRO	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	lethal minimum value of leaf area ratio	(m^2/kg)
XUNIF_CNA_NITRO CFNAM_CNA_NITRO CFTYP_CNA_NITRO	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	nitrogen concentration of active biomass	(kg/kg)

TEB scheme

Over urban areas, all surface parameters have to be specified by the user in namelist **NAM_DATA_TEB**.

Fortran name	Fortran type	values	default	description	unit
XUNIF_BLD CFNAM_BLD CFTYP_BLD	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	fraction of buildings	(-)
XUNIF_BLD_HEIGHT CFNAM_BLD_HEIGHT CFTYP_BLD_HEIGHT	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	buildings height	(m)
XUNIF_WALL_O_HOR CFNAM_WALL_O_HOR CFTYP_WALL_O_HOR	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	wall surf. / hor. surf.	(-)
XUNIF_Z0_TOWN	real		none	roughness length for momentum	(m)
CFNAM_Z0_TOWN CFTYP_Z0_TOWN	character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	, , none		
XUNIF_ALB_ROOF CFNAM_ALB_ROOF CFTYP_ALB_ROOF	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	roof albedo	(-)
XUNIF_EMIS_ROOF CFNAM_EMIS_ROOF CFTYP_EMIS_ROOF	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	roof emissivity	(-)
XUNIF_HC_ROOF CFNAM_HC_ROOF CFTYP_HC_ROOF	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	roof layers heat capacity	(J/K/m ³)
XUNIF_TC_ROOF CFNAM_TC_ROOF CFTYP_TC_ROOF	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	roof layers thermal conductivity	(W/K/m)
XUNIF_D_ROOF CFNAM_D_ROOF CFTYP_D_ROOF	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	roof layers depth	(m)
XUNIF_ALB_ROAD CFNAM_ALB_ROAD CFTYP_ALB_ROAD	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	road albedo	(-)
XUNIF_EMIS_ROAD CFNAM_EMIS_ROAD CFTYP_EMIS_ROAD	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	road emissivity	(-)
XUNIF_HC_ROAD CFNAM_HC_ROAD CFTYP_HC_ROAD	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	road layers heat capacity	(J/K/m ³)
XUNIF_TC_ROAD CFNAM_TC_ROAD CFTYP_TC_ROAD	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	road layers thermal conductivity	(W/K/m)
XUNIF_D_ROAD CFNAM_D_ROAD CFTYP_D_ROAD	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none , , none	road layers depth	(m)

Fortran name	Fortran type	values	default	description	unit
XUNIF_ALB_WALL	real		none	wall albedo	(-)
CFNAM_ALB_WALL	character (LEN=28)		, ,		
CFTYP_ALB_WALL	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			
XUNIF_EMIS_WALL	real		none	wall emissivity	(-)
CFNAM_EMIS_WALL	character (LEN=28)		, ,		
CFTYP_EMIS_WALL	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			
XUNIF_HC_WALL	real		none	wall layers heat capacity	(J/K/m ³)
CFNAM_HC_WALL	character (LEN=28)		, ,		
CFTYP_HC_WALL	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			
XUNIF_TC_WALL	real		none	wall layers thermal conductivity	(W/K/m)
CFNAM_TC_WALL	character (LEN=28)		, ,		
CFTYP_TC_WALL	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			
XUNIF_D_WALL	real		none	wall layers depth	(m)
CFNAM_D_WALL	character (LEN=28)		, ,		
CFTYP_D_WALL	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			
XUNIF_H_TRAFFIC	real		none	anthropogenic sensible heat fluxes due to traffic	(W/m ²)
CFNAM_H_TRAFFIC	character (LEN=28)		, ,		
CFTYP_H_TRAFFIC	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			
XUNIF_LE_TRAFFIC	real		none	anthropogenic latent heat fluxes due to traffic	(W/m ²)
CFNAM_LE_TRAFFIC	character (LEN=28)		, ,		
CFTYP_LE_TRAFFIC	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			
XUNIF_H_INDUSTRIES	real		none	anthropogenic sensible heat fluxes due to factories	(W/m ²)
CFNAM_H_INDUSTRIES	character (LEN=28)		, ,		
CFTYP_H_INDUSTRIES	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			
XUNIF_LE_INDUSTRIES	real		none	anthropogenic latent heat fluxes due to factories	(W/m ²)
CFNAM_LE_INDUSTRIES	character (LEN=28)		, ,		
CFTYP_LE_INDUSTRIES	character (LEN=6)	'DIRECT', 'BINLLF'	none		
		'BINLLV', 'ASCLLV'			

SEAFLUX scheme

Treat SST as a forcing variable. For that purpose, several SST files at a given time are required and namelist **NAM_DATA_SEAFLUX** should be filled.

Fortran name	Fortran type	values	default	description	unit
LSST_DATA	logical		none	flag to activate this option	(-)
NTIME	integer	12	12	number of SST data	(-)
CFNAM_SST	character (LEN=28)		' '		
CFTYP_SST	character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	none		
NYEAR_SST	integer				
NMONTH_SST	integer				
NDAY_SST	integer				
XTIME_SST	real				

- LSST_DATA : flag to initialize SST from a climatology
- NTIME : number of SST input files
- CFNAM_SST : SST data file name
- CFTYP_SST : type of SST data file ('DIRECT', 'BINLLF', 'BINLLV', 'ASCLLV')
- NYEAR_SST : year of SST data file
- NMONTH_SST : month of SST data file
- NDAY_SST : day of SST data file
- XTIME_SST : time in seconds of SST data file

How to initialise SST from external files : an example with 3 SST input files (lat, lon, value type).

```

&NAM_DATA_SEAFLUX      NTIME = 3 , LSST_DATA = T ,
                        CFNAM_SST (1)      = 'sst_1.dat'           , CFTYP_SST (1)      = 'ASCLLV',
                        CFNAM_SST (2)      = 'sst_2.dat'           , CFTYP_SST (2)      = 'ASCLLV',
                        CFNAM_SST (3)      = 'sst_3.dat'           , CFTYP_SST (3)      = 'ASCLLV',
                        NYEAR_SST(1)=1985,  NMONTH_SST(1)=12,  NDAY_SST(1)=31 , XTIME_SST(1)=64800.,
                        NYEAR_SST(2)=1986,  NMONTH_SST(2)=1 ,  NDAY_SST(2)=1  , XTIME_SST(2)=43200.,
                        NYEAR_SST(3)=1986,  NMONTH_SST(3)=1 ,  NDAY_SST(3)=2  , XTIME_SST(3)=0.

```

/

FLAKE scheme

Over lakes, if one wants to use Flake scheme, some parameters have to be specified by the user in the namelist **NAM_DATA_FLAKE**.

Fortran name	Fortran type	values	default	description	unit
XUNIF_WATER_DEPTH YWATER_DEPTH YWATER_DEPTHFILETYPE	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	20. ' ' none	Lake depth filename	(<i>m</i>)
XUNIF_WATER_FETCH YWATER_FETCH YWATER_FETCHFILETYPE	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	1000. ' ' none	wind fetch filename	(<i>m</i>)
XUNIF_T_BS YT_BS YT_BSFILETYPE	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	286. ' ' none	temperature at the outer edge of the thermally active layer of the bottom sediments filename	(<i>K</i>)
XUNIF_DEPTH_BS YDEPTH_BS YDEPTH_BSFILETYPE	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	1. ' ' none	depth of the sediments layer filename	(<i>m</i>)
XUNIF_EXTCOEFF_WATER YEXTCOEFF_WATER YEXTCOEFF_WATERFILETYPE	real character (LEN=28) character (LEN=6)	'DIRECT', 'BINLLF' 'BINLLV', 'ASCLLV'	3. ' ' none	extinction coefficient of solar radiation in water filename	(<i>m</i> ⁻¹)

- XUNIF_xxx : uniform prescribed value of parameter xxx. If XUNIF_xxx is set, file CFNAM_xxx is not used.
- CFNAM_xxx : data file name associated to parameter xxx. If XUNIF_xxx is set, file CFNAM_xxx is not used.
- CFTYP_xxx : type of sea data file ('DIRECT', 'BINLLF', 'BINLLV', 'ASCLLV')

8.2 Choice of the surface schemes

You must first choose the surface schemes you will use. It is not possible, once chosen, to modify the surface schemes in the later steps (**PREP**, **running of the schemes**, **DIAG**).

Depending on the schemes you use, some additional physiographic fields will be computed if they are needed for the surface scheme chosen. For example, the ISBA scheme (used for vegetation and soil) needs the fractions of clay and sand.

Namelist NAM_PGD_SCHEMES

This namelist defines the four schemes that will be used, one for each type of surface (sea, inland water, town, vegetation).

Fortran name	Fortran type	values	default value
CNATURE	string of 6 characters	"NONE " , "FLUX " , "TSZ0 " , "ISBA "	"ISBA "
CSEA	string of 6 characters	"NONE " , "FLUX " , "SEAFIX"	"SEAFIX"
CWATER	string of 6 characters	"NONE " , "FLUX " , "WATFIX" , "FLAKE "	"WATFIX"
CTOWN	string of 6 characters	"NONE " , "FLUX " , "TEB "	"TEB "

- CNATURE : scheme used for vegetation and natural soil covers . The different possibilities are :
 1. "NONE " : no scheme used. No fluxes will be computed at the surface.
 2. "FLUX " : ideal fluxes are prescribed. They have to be set in the fortran routine *init_ideal_flux.f90*.
 3. "TSZ0 " : In this scheme, the fluxes are computed according to the ISBA physics, but the surface characteristics (temperature, humidity, etc...) remain constant with time.
 4. "ISBA " : this is the full ISBA scheme (Noilhan and Planton 1989), with all options developed since this initial paper.
- CSEA : scheme used for sea and ocean . The different possibilities are :
 1. "NONE " : no scheme used. No fluxes will be computed at the surface.
 2. "FLUX " : ideal fluxes are prescribed. They have to be set in the fortran routine *init_ideal_flux.f90*.
 3. "SEAFIX" : this is a relatively simple scheme, using the Charnock formula.
- CWATER : scheme used for inland water . The different possibilities are :
 1. "NONE " : no scheme used. No fluxes will be computed at the surface.
 2. "FLUX " : ideal fluxes are prescribed. They have to be set in the fortran routine *init_ideal_flux.f90*.
 3. "WATFIX" : this is a relatively simple scheme, using the Charnock formula.
 4. "FLAKE " : this is lake scheme from Mironov, 2005.
- CTOWN : scheme used for towns . The different possibilities are :
 1. "NONE " : no scheme used. No fluxes will be computed at the surface.
 2. "FLUX " : ideal fluxes are prescribed. They have to be set in the fortran routine *init_ideal_flux.f90*.
 3. "TEB " : this is the Town Energy Balance scheme (Masson 2000), with all the subsequent ameliorations of the scheme.

8.3 Definition of the grid

Note that all the namelists presented in this section are ignored if the grid is imposed, in the fortran code, from an atmospheric model. This is the case when one already have defined the atmospheric grid and one want to be sure that the surface has the same grid. For example, this is what happens in the MESONH program PREP.IDEAL.CASE (when no physiographic surface file is used).

If you are in this case, ignore all the namelists presented in this section, and only the namelists for cover and the following ones, have to be used.

8.3.1 Choice of the grid type

Namelist NAM_PGD_GRID

This namelist defines the grid type, either specified or from an existing surface file

Fortran name	Fortran type	default value
CGRID	string of 10 characters	"CONF PROJ "
YINIFILE	string of 28 characters	none
YFILETYPE	string of 6 characters	none

- CGRID : type of grid and projection . It is used **only** if a file is not prescribed (see below). The different grid possibilities are :
 1. "GAUSS " : this grid is a gaussian grid (global grid, that may be stretched, rotated, ...)
 2. "CONF PROJ " : this grid is a regular grid (in meters in x and y perpendicular directions) on conformal projection plan (Mercator, Lambert or polar stereographic).
 3. "CARTESIAN " : this grid is a regular grid (in meters in x and y perpendicular directions), with no reference to real geographical coordinates.
 4. "LONLAT REG" : this grid is defined as a regular latitude - longitude grid.
 5. "IGN " : this grid type contains all IGN (French National Geographical Institute) possible Lambert projections
 6. "NONE " : this grid is not regular. Only the number of points and the size of each grid mesh is prescribed. There is no positioning of each point compared to any other.
- YINIFILE : name of the file used to define the grid. It is possible to define the grid as a subgrid of a previously created file. This is currently possible only for files that have a "CONF PROJ " or "CARTESIAN " grid type. The exact definition of the subgrid grid chosen is prescribed in a namelist (described below), depending on the type of grid available in the file chosen. The use of a file has priority on the CGRID type.
- YFILETYPE : type of the YINIFILE file, if the latter is provided. YFILETYPE must be given. The following values are currently usable :
 - "MESONH" : the file type is a MESONH file.

8.3.2 Conformal projection grids (Mercator, Lambert, Polar stereographic)

Namelist NAM_CONF_PROJ

This namelist defines the projection in case CGRID="CONF PROJ "

Fortran name	Fortran type	default value
XLAT0	real	none
XLON0	real	none
XRPK	real	none
XBETA	real	none

- XLAT0 : reference latitude for conformal projection (real, decimal degrees)
- XLON0 : reference longitude for conformal projection (real, decimal degrees)
- XRPK : cone factor for the projection (real) :
 - XRPK=1 : polar stereographic projection from south pole
 - 1>XRPK>0 : Lambert projection from south pole
 - XRPK=0 : Mercator projection from earth center
 - -1<XRPK<0 : Lambert projection from north pole
 - XRPK=-1 : polar stereographic projection from north pole
- XBETA : rotation angle of the simulation domain around the reference longitude (real)

Namelist NAM_CONF_PROJ_GRID

This namelists defines the horizontal domain in case CGRID="CONF PROJ ".

Fortran name	Fortran type
XLATCEN	real
XLONCEN	real
NIMAX	integer
NJMAX	integer
DX	real
DY	real

- XLATCEN : latitude of the point of the center of the domain (real, decimal degrees)
- XLONCEN : longitude of the point of the center of the domain (real, decimal degrees)
- NIMAX : number of surface points of the grid in direction x .
- NJMAX : number of surface points of the grid in direction y .
- DX : grid mesh size on the conformal plane in x direction (real, meters).
- DY : grid mesh size on the conformal plane in y direction (real, meters).

Namelist NAM_INIFILE_CONF_PROJ

This namelists defines the horizontal domain from an existing surface file in which grid type is "CONF PROJ ". If nothing is set in the namelist, a grid identical as the one in the file is chosen.

Fortran name	Fortran type	default value
IXOR	integer	1
IYOR	integer	1
IXSIZE	integer	YINIFILE size
IYSIZE	integer	YINIFILE size
IDXRATIO	integer	1
IDYRATIO	integer	1

- IXOR : first point I index, according to the YINIFILE grid, left to and out of the new physical domain.
- IYOR : first point J index, according to the YINIFILE grid, under and out of the new physical domain.
- IXSIZE : number of grid points in I direction, according to YINIFILE grid, recovered by the new domain. If to be used in MESONH, it must only be factor of 2,3 or 5.
- IYSIZE : number of grid points in J direction, according to YINIFILE grid, recovered by the new domain. If to be used in MESONH, it must only be factor of 2,3 or 5.
- IDXRATIO : resolution factor in I direction between the YINIFILE grid and the new grid. If to be used in MESONH, it must only be factor of 2,3 or 5.
- IDYRATIO : resolution factor in J direction between the YINIFILE grid and the new grid. If to be used in MESONH, it must only be factor of 2,3 or 5.

8.3.3 Cartesian grids

Namelist NAM_CARTESIAN

This namelist defines the projection in case CGRID="CARTESIAN "

Fortran name	Fortran type	default value
XLAT0	real	none
XLON0	real	none
NIMAX	integer	none
NJMAX	integer	none
XDX	real	none
XDY	real	none

- XLAT0 : reference latitude (real, decimal degrees)
- XLON0 : reference longitude (real, decimal degrees)
- NIMAX : number of surface points of the grid in direction x .
- NJMAX : number of surface points of the grid in direction y .
- XDX : grid mesh size on the conformal plane in x direction (real, meters).
- XDY : grid mesh size on the conformal plane in y direction (real, meters).

Namelist NAM_INIFILE_CARTESIAN

This namelists defines the horizontal domain from an existing surface file in which grid type is "CARTESIAN ". If nothing is set in the namelist, a grid identical as the one in the file is chosen.

Fortran name	Fortran type	default value
IXOR	integer	1
IYOR	integer	1
IXSIZE	integer	YINIFILE size
IYSIZE	integer	YINIFILE size
IDXRATIO	integer	1
IDYRATIO	integer	1

- IXOR : first point I index, according to the YINIFILE grid, left to and out of the new physical domain.
- IYOR : first point J index, according to the YINIFILE grid, under and out of the new physical domain.
- IXSIZE : number of grid points in I direction, according to YINIFILE grid, recovered by the new domain. If to be used in MESONH, it must only be factor of 2,3 or 5.
- IYSIZE : number of grid points in J direction, according to YINIFILE grid, recovered by the new domain. If to be used in MESONH, it must only be factor of 2,3 or 5.
- IDXRATIO : resolution factor in I direction between the YINIFILE grid and the new grid. If to be used in MESONH, it must only be factor of 2,3 or 5.
- IDYRATIO : resolution factor in J direction between the YINIFILE grid and the new grid. If to be used in MESONH, it must only be factor of 2,3 or 5.

8.3.4 Regular longitude-latitude grids

Namelist NAM_LONLAT_REG

This namelist defines the projection in case CGRID="LONLAT REG"

Fortran name	Fortran type	default value
XLONMIN	real	none
XLONMAX	real	none
XLATMIN	real	none
XLATMAX	real	none
NLON	integer	none
NLAT	integer	none

- XLONMIN : minimum longitude covered by the grid, i.e. corresponding to the west border of the domain (real, decimal degrees). XLONMIN must be smaller than XLONMAX, but no more than 360 smaller.
- XLONMAX : maximum longitude covered by the grid, i.e. corresponding to the east border of the domain (real, decimal degrees). XLONMAX must be larger than XLONMIN, but no more than 360 larger.
- XLATMIN : minimum latitude covered by the grid, i.e. corresponding to the south border of the domain (real, decimal degrees). XLATMIN must be between -90 and +90, and smaller than XLATMAX.
- XLATMAX : maximum longitude covered by the grid, i.e. corresponding to the 'right' border of the domain (real, decimal degrees). XLATMAX must be between -90 and +90, and larger than XLATMIN.
- NLON : number of surface points in the longitude direction.
- NLAT : number of surface points in the latitude direction.

8.3.5 Regular Lambert grids

Namelist NAM_IGN

This namelist defines the projection in case CGRID="IGN "

Fortran name	Fortran type	values	default value
CLAMBERT	character (len=3)	'L1 ', 'L2 ', 'L3 ', 'L4 ', 'L2E', 'L93'	none
NPOINTS	integer		none
XX	real		none
XY	real		none
XDX	real		none
XDY	real		none

- CLAMBERT : type of Lambert prjection
 - "L1 " : Lambert I
 - "L2 " : Lambert II
 - "L3 " : Lambert III
 - "L4 " : Lambert IV
 - "L2E" : Extended Lambert II
 - "L93" : Lambert 93
- NPOINTS : number of grid points defining the grid
- XX : X coordinate of grid mesh center
- YY : Y coordinate of grid mesh center
- XDX : grid mesh size on the conformal plane in x direction (real, meters).
- XDY : grid mesh size on the conformal plane in y direction (real, meters).

8.3.6 Gaussian grids

These namelists define the projection in case CGRID="GAUSS "

Namelist NAMDIM

Fortran name	Fortran type	default value
NDGLG	integer	none

Namelist NAMRGRI

Fortran name	Fortran type	default value
NRGRI	integer	none

Namelist NAMGEM

Fortran name	Fortran type	default value
RMUCEN	real	none
RLOCEN	real	none
RSTRET	real	none

- NDGLG : number of pseudo-latitudes
- NRGRI : number of pseudo-longitudes on each pseudo-latitude circle starting from the rotated pole
- RMUCEN : sine of the latitude of the rotated pole
- RLOCEN : longitude of the rotated pole (radian)
- RSTRET : stretching factor (must be greater than or equal to 1)

8.4 Land cover fractions

Namelist NAM_COVER

This namelist gives the information to compute the surface cover fractions.

Fortran name	Fortran type	values	default value
XUNIF_COVER	array of 255 reals	≥ 0 and $\sum_{i=1}^{255} \text{XUNIF\_COVER}(i) = 1$	none
YCOVER	character (LEN=28)		' '
YFILETYPE	character (LEN=6)	'DIRECT', 'BINLLF', 'BINLLV', 'ASCLLV'	none
XRM_COVER	real	≥ 0	10^{-6}
LRM_TOWN	logical		.FALSE.

- XUNIF_COVER : specified values for uniform cover fractions. For each index i between 1 and 255, XUNIF_COVER(i) is the fraction of the i^{th} ecosystem of ecoclimap. The same fraction of each ecosystem is set to all points of the grid. The sum of all ecosystem fractions must be equal to one : $\sum_{i=1}^{255} \text{XUNIF\_COVER}(i) = 1$.
If XUNIF_COVER is set, it has priority on the use of an ecosystem file (see next item : YCOVER). In the case of grid without any reference to geographical coordinates ("CARTESIAN " or "NONE "), XUNIF_COVER **must** be set.
- YCOVER : ecoclimap data file name. It is used only if XUNIF_COVER is not set.
- YFILETYPE : type of YCOVER file ('DIRECT', 'BINLLV', 'BINLLF', 'ASCLLV').
- XRM_COVER : for each point, all fractions of ecosystems that are below XRM_COVER are removed (i.e. set to zero), and the corresponding area fractions are distributed among the remaining ecosystem fractions. Whatever the value of XRM_COVER, at least one ecosystem remains for each grid point.
- LRM_TOWN : if .TRUE., all ecosystems containing town fractions are removed and replaced by the ecosystem corresponding to rocks.

8.5 Orography and subgrid orography

Namelist NAM_ZS

This namelist defines the orography file and orographic treatment to be done.

Fortran name	Fortran type	values	default value
XUNIF_ZS	real		none
YZS	character (LEN=28)		' ' (default orography is 0.)
YFILETYPE	character (LEN=6)	'DIRECT', 'BINLLF'	none
		'BINLLV', 'ASCLLV'	
COROGTYPE	character (LEN=3)	'AVG', 'ENV', 'SIL '	'ENV'
XENV	real		0.
NZSFILTER	integer		1

- XUNIF_ZS : uniform value of orography imposed on all points (real,meters). If XUNIF_ZS is set, file YZS is not used.
- YZS : data file name. If XUNIF_ZS is set, file YZS is not used. If neither XUNIF_ZS and YZS is set, then orography is set to zero.
- YFILETYPE : type of data file ('DIRECT', 'BINLLF', 'BINLLV', 'ASCLLV')
- COROGTYPE : type of orography (string of 3 characters) :
 - 'AVG' : mean orography $\overline{z_s}$.
 - 'ENV' : envelope relief, defined from mean orography and the subgrid orography standard deviation as $\overline{z_s} + XENV * \sigma_{z_s}$.
 - 'SIL' : silhouette relief, defined as the mean of the two subgrid silhouettes in directions x and y (if two main directions can be defined for the grid chosen).
- XENV : enhance factor in envelope orography definition (real).
- NZSFILTER : number of iterations of the spatial filter applied to smooth the orography (integer, 1 iteration removes the $2\Delta x$ signal, 50% of the $4\Delta x$ signal, 25% of the $6\Delta x$ signal, etc¹...).

¹The amplitude of the filtered signal for each wavelength $\lambda\Delta x$ is $\frac{1}{2}(\cos(2\pi/\lambda) + 1)$.

8.6 Namelist for ISBA scheme

Namelist NAM_ISBA

Fortran name	Fortran type	values	default value
NPATCH	integer	between 1 and 12	1
CISBA	character (LEN=3)	'2-L', '3-L', 'DIF'	'3-L'
CPHOTO	string of 3 characters	'NON', 'AGS', 'LAI', 'AST', 'LST', 'NIT'	'NON'
NGROUND_LAYER	integer	> 0	3
XUNIF_CLAY	real	between 0 and 1	none
YCLAY	character (LEN=28)		' '
YCLAYFILETYPE	character (LEN=6)	'DIRECT', 'BINLLF'	none
XUNIF_SAND	real	'BINLLV', 'ASCLLV'	none
YSAND	character (LEN=28)	between 0 and 1	' '
YSANDFILETYPE	character (LEN=6)	'DIRECT', 'BINLLF'	none
		'BINLLV', 'ASCLLV'	
XUNIF_RUNOFFB	real	between 0 and 1	0.5

- NPATCH : number of patches used in ISBA. One patch corresponds to aggregated parameters. 12 patches correspond to separate energy budgets for all vegetation types present in ISBA. 3 patches correspond to bare soil types, low vegetation, trees. If CPHOTO equals 'NON' any number of patches between 1 and 12 is possible, for the other values of CPHOTO, 12 patches are required. The order and the signification of each patch is the following :
 - 1 : bare ground
 - 2 : rocks
 - 3 : permanent snow
 - 4 : deciduous forest
 - 5 : conifer forest
 - 6 : evergreen broadleaf trees
 - 7 : C3 crops
 - 8 : C4 crops
 - 9 : irrigated crops
 - 10 : grassland (C3)
 - 11 : tropical grassland (C4)
 - 12 : garden and parks
- CISBA : type of soil discretization and physics in ISBA :
 - '2-L' : force-restore method with 2 layers for hydrology
 - '3-L' : force-restore method with 3 layers for hydrology
 - 'DIF' : diffusion layer, with any number of layers
- CPHOTO : type of photosynthesis physics. The following options are currently available :
 - "NON" : none is used, Jarvis formula is used for plant transpiration.
 - "AGS" : ISBA-AGS, without evolving Leaf Area Index
 - "LAI" : ISBA-AGS, with evolving Leaf Area Index
 - "AST" : ISBA-AGS with offensive/defensive stress, without evolving Leaf Area Index
 - "LST" : ISBA-AGS with offensive/defensive stress, with evolving Leaf Area Index
 - "NIT" : ISBA-AGS with nitrogen, with evolving Leaf Area Index
- NGROUND_LAYER : number of soil layer used in case of diffusion physics in the soil (CISBA = 'DIF').
- XUNIF_CLAY : uniform prescribed value of clay fraction. If XUNIF_CLAY is set, file YCLAY is not used.
- YCLAY : clay fraction data file name. If XUNIF_CLAY is set, file YCLAY is not used.
- YCLAYFILETYPE : type of clay data file ('DIRECT', 'BINLLF', 'BINLLV', 'ASCLLV')
- XUNIF_SAND : uniform prescribed value of sand fraction. If XUNIF_SAND is set, file YSAND is not used.
- YSAND : sand fraction data file name. If XUNIF_SAND is set, file YSAND is not used.
- YSANDFILETYPE : type of sand data file ('DIRECT', 'BINLLF', 'BINLLV', 'ASCLLV')
- XUNIF_RUNOFFB : uniform prescribed value of subgrid runoff coefficient.

8.7 Namelist to add user's own fields

Namelist NAM_DUMMY_PGD

This namelist allows to incorporate into the physiographic file any surface field.

You can treat up to 999 such fields. These fields will be written on all the files you will use later(after prognostic fields initialization, or during and after run, etc...). Their name in the files are 'DUMMY_GRnnn', where nnn goes from 001 to 999.

During the execution of the programs, these fields are stored in the XDUMMY_FIELDS(:, :) (first dimension : spatial dimension, second dimension : total number of fields), in the module MODD_DUMMY_SURF_FIELD\$. You must modify the fortran source, where you want to use them.

Fortran name	Fortran type	default value
NDUMMY_PGD_NBR	integer	0
CDUMMY_PGD_NAME(:)	1000 * character (LEN=20)	1000 * ' '
CDUMMY_PGD_FILE(:)	1000 * character (LEN=28)	1000 * ' '
CDUMMY_PGD_COMMENT(:)	1000 * character (LEN=40)	1000 * ' '
CDUMMY_PGD_FILETYPE(:)	1000 * character (LEN=6)	1000 * ' '
CDUMMY_PGD_AREA(:)	1000 * character (LEN=3)	1000 * 'ALL'
CDUMMY_PGD_ATYPE(:)	1000 * character (LEN=3)	1000 * 'ARI'

Only the first NDUMMY_PGD_NBR values in these arrays are meaningful.

- NDUMMY_PGD_NBR : number of dummy fields.
- CDUMMY_PGD_NAME(:) : list of the dummy fields you want to initialize with your own data. You can give any name you want. This is a way to describe what is the field. This information is not used by the program. It is just written in the FM files.
- CDUMMY_PGD_FILE(:) : list of the names of the files containing the data for the fields you have specified in CDUMMY_PGD_NAME(:).
- CDUMMY_PGD_FILETYPE(:) : list of the types of the files containing the data for the fields you have specified in CDUMMY_PGD_NAME(:) ('DIRECT', 'LATLON', 'BINLLF', 'BINLLV', 'ASCLLV').
- CDUMMY_PGD_AREA(:) : area of meaningfulness of the fields you have specified in CDUMMY_PGD_NAME(:) ('ALL', 'NAT', 'TWN', 'SEA', 'WAT', 'LAN', respectively for everywhere, natural areas, town areas, sea, inland waters, land = natural cover + town). For example, oceanic emission of DNS is relevant on 'SEA'.
- CDUMMY_PGD_ATYPE(:) : type of averaging (during **PGD** for the fields you have specified in CDUMMY_PGD_NAME(:) ('ARI', 'INV', 'LOG', respectively for arithmetic, inverse and logarithmic averaging).

8.8 Namelist for chemistry anthropogenic emissions

Namelist NAM_CH_EMIS_PGD

This namelist is used to initialize chemistry components emissions.

You can treat up to 999 such fields. These fields will be written on all the files you will use later (after prognostic fields initialization, or during and after run, etc...). Their name in the files are 'EMIS_GRnnn', where nnn goes from 001 to 999.

During the execution of the programs, these fields are stored in the XEMIS_GR_FIELDS(:, :) (first dimension : spatial dimension, second dimension : total number of fields), in the module MODD_EMIS_GR_FIELD\$. The temporal evolution, the aggregation of prescribed emissions and the link with the corresponding chemical prognostic variables are handled by the subroutine CH_EMISSION_FLUXn.f90

Fortran name	Fortran type	default value
NEMIS_PGD_NBR	integer	0
CEMIS_PGD_NAME(:)	1000 * character (LEN=20)	1000 * ' '
CEMIS_PGD_FILE(:)	1000 * character (LEN=28)	1000 * ' '
CEMIS_PGD_COMMENT(:)	1000 * character (LEN=40)	1000 * ' '
NEMIS_PGD_TIME	integer	0
CEMIS_PGD_FILETYPE(:)	1000 * character (LEN=6)	1000 * 'DIRECT'
CEMIS_PGD_AREA(:)	1000 * character (LEN=3)	1000 * 'ALL'
CEMIS_PGD_ATYPE(:)	1000 * character (LEN=3)	1000 * 'ARI'

Only the first NEMIS_PGD_NBR values in these arrays are meaningful.

- NEMIS_PGD_NBR : number of dummy fields.
- CEMIS_PGD_NAME(:) : list of the dummy fields you want to initialize with your own data. You can give any name you want. This is a way to describe what is the field. This information is not used by the program. It is just written in the FM files.
- CEMIS_PGD_FILE(:) : list of the names of the files containing the data for the fields you have specified in CEMIS_PGD_NAME(:).
- CEMIS_PGD_COMMENT(:) : list of the comments associated to each emission field.
- NEMIS_PGD_TIME(:) : list of the time of the files containing the data for the fields you have specified in CEMIS_PGD_NAME(:).
- CEMIS_PGD_FILETYPE(:) : list of the types of the files containing the data for the fields you have specified in CEMIS_PGD_NAME(:) ('DIRECT', 'BINLLF', 'BINLLV', 'ASCLLV').
- CEMIS_PGD_AREA(:) : area of meaningfulness of the fields you have specified in CEMIS_PGD_NAME(:) ('ALL', 'NAT', 'TWN', 'SEA', 'WAT', 'LAN', respectively for everywhere, natural areas, town areas, sea, inland waters, land = natural cover + town). For example, oceanic emission of DNS is relevant on 'SEA'.
- CEMIS_PGD_ATYPE(:) : type of averaging (during **PGD** for the fields you have specified in CEMIS_PGD_NAME(:) ('ARI', 'INV', 'LOG', respectively for arithmetic, inverse and logarithmic averaging).

Example :

```
&NAM_CH_EMIS_PGD NEMIS_PGD_NBR = 2,  
  CEMIS_PGD_NAME(1)='COE',  
  NEMIS_PGD_TIME(1)=0,  
  CEMIS_PGD_COMMENT(1)='CO_00h00',  
  CEMIS_PGD_AREA(1)='LAN',  
  CEMIS_PGD_ATYPE(1)='ARI',  
  CEMIS_PGD_FILE(1)='co_00.asc',  
  CEMIS_PGD_FILETYPE(1)='ASCLLV',  
  CEMIS_PGD_NAME(2)='COE',  
  NEMIS_PGD_TIME(2)=43200,  
  CEMIS_PGD_COMMENT(2)='CO_12h00',  
  CEMIS_PGD_AREA(2)='LAN',  
  CEMIS_PGD_ATYPE(2)='ARI',  
  CEMIS_PGD_FILE(2)='co_12.asc',  
  CEMIS_PGD_FILETYPE(2)='ASCLLV',  
  CEMIS_PGD_NAME(3)='DMSE',  
  NEMIS_PGD_TIME(3)=0,  
  CEMIS_PGD_COMMENT(3)='dms_cte',  
  CEMIS_PGD_AREA(3)='SEA',  
  CEMIS_PGD_ATYPE(3)='ARI',  
  CEMIS_PGD_FILE(3)='dms.asc',  
  CEMIS_PGD_FILETYPE(3)='ASCLLV' /
```


Chapitre 9

Initialization of the prognostic fields

9.1 Overview of fields computation : PREP

The prognostic fields (temperature, humidity, ice, snow, etc...) are averaged or interpolated on the specified grid by the program **PREP**. They are stored in the surface file. The computation is done separately for each surface scheme.

During the **PREP** facility :

1. You initializes the date of the surface
2. You initializes the prognostic variables of the chosen sea scheme
3. You initializes the prognostic variables of the chosen lake scheme
4. You initializes the prognostic variables of the chosen vegetation scheme
5. You initializes the prognostic variables of the chosen town scheme

Here are presented the initialization procedures for the schemes that need such information (for example, scheme "IDEAL " does not need any information here, but modification of the code source *init_ideal_flux.f90*).

Note that for each scheme, and for some for each variable of the scheme, it is possible to initialize the prognostic fields either form an operationnal or research model, or using prescribed (usually uniform) fields.

9.2 Date initialization and default input data file for all schemes

Namelist NAM_PREP_SURF_ATM

This namelist information is used to (possibly) :

- initialize the date of all surface schemes. The namelist information is used only if no input data file is used, either from namelist or by fortran code (as in MESONH programs). If a file is used, the date is read in it.
- define the default file in which each scheme can read the needed data (e.g. temperature).

Note that, all the information given in this namelist can be erased for each scheme by the namelist corresponding to this scheme, as the information in the scheme namelists have priority on namelist NAM_PREP_SURF_ATM.

Fortran name	Fortran type	values	default value
CFILE	string of 28 characters		- atmospheric file used in the program calling the surface facilities, if any - none otherwise
CFILETYPE	string of 6 characters	'MESONH', 'GRIB '	- type of the atmospheric file, if any - none otherwise
NYEAR	integer		none
NMONTH	integer		none
NDAY	integer		none
XTIME	real		none

- CFILE : name of the file used to define
 1. the date.
 2. the file in which to read the needed data (e.g. temperature).

The use of a file or prescribed value in each scheme namelist has priority on the data in CFILE file of namelist NAM_PREP_SURF_ATM.

- CFILETYPE : type of the CFILE file, if the latter is provided. CTYPE must then be given. The following values are currently usable :
 - "MESONH" : the file type is a MESONH file.
 - "GRIB " : the file type is a GRIB file, coming from any of these models :
 1. "ECMWF " : european center forecast model
 2. "ARPEGE" : Arpege french forecast model
 3. "ALADIN" : Aladin french forecast local model
 4. "MOCAGE" : Mocage french research chemistry model
- NYEAR : year of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read).
- NMONTH : month of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read).
- NDAY : day of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read).
- XTIME : time from midnight of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read). (seconds).

9.3 Sea scheme "SEAFLEX"

Namelist NAM_PREP_SEAFLEX

This namelist information is used to initialize the "SEAFLEX" sea scheme temperature.

Fortran name	Fortran type	values	default value
XSST_UNIF	real		none
CFILE_SEAFLEX	string of 28 characters		CFILE in NAM_PREP_SURF_ATM
CTYPE	string of 6 characters	'MESONH', 'GRIB '	CFILETYPE in NAM_PREP_SURF_ATM
NYEAR	integer		none
NMONTH	integer		none
NDAY	integer		none
XTIME	real		none
LSEA_SBL	logical		F

- XSST_UNIF : uniform prescribed value of Sea Surface Temperature. This prescribed value, if defined, has priority on the use of CFILE_SEAFLEX data.
- CFILE_SEAFLEX : name of the file used to define the Sea surface Temperature. **The use of a file or prescribed value XSST_UNIF has priority on the data in CFILE_SEAFLEX file.**
- CTYPE : type of the CFILE_SEAFLEX file, if the latter is provided. CTYPE must then be given. The following values are currently usable :
 - "MESONH" : the file type is a MESONH file.
 - "GRIB " : the file type is a GRIB file, coming from any of these models :
 1. "ECMWF " : european center forecast model
 2. "ARPEGE" : Arpege french forecast model
 3. "ALADIN" : Aladin french forecast local model
 4. "MOCAGE" : Mocage french research chemistry model
- NYEAR : year of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read).
- NMONTH : month of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read).
- NDAY : day of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read).
- XTIME : time from midnight of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read). (seconds).
- LSEA_SBL : activates surface boundary multi layer scheme over sea.

9.4 Lake scheme "WATFLX"

Namelist NAM_PREP_WATFLUX

This namelist information is used to initialize the "WATFLX" sea scheme temperature.

Fortran name	Fortran type	values	default value
XTS_WATER_UNIF	real		none
CFILE_WATFLX	string of 28 characters		CFILE in NAM_PREP_SURF_ATM
CTYPE	string of 6 characters	'MESONH', 'GRIB '	CFILETYPE in NAM_PREP_SURF_ATM
NYEAR	integer		none
NMONTH	integer		none
NDAY	integer		none
XTIME	real		none
LWAT_SBL	logical		F

- XTS_WATER_UNIF : uniform prescribed value of water surface temperature supposed at an altitude of 0m (mean sea level altitude). The temperature is then modified for each point depending on its altitude, following a uniform vertical gradient of -6.5 Kkm^{-1} . This prescribed value, if defined, has priority on the use of CFILE_WATFLX data.
- CFILE_WATFLX : name of the file used to define the Sea surface Temperature. **The use of a file or prescribed value XTS_WATER_UNIF has priority on the data in CFILE_WATFLX file.**
- CTYPE : type of the CFILE_WATFLX file, if the latter is provided. CTYPE must then be given. The following values are currently usable :
 - "MESONH" : the file type is a MESONH file.
 - "GRIB " : the file type is a GRIB file, coming from any of these models :
 1. "ECMWF " : european center forecast model
 2. "ARPEGE" : Arpege french forecast model
 3. "ALADIN" : Aladin french forecast local model
 4. "MOCAGE" : Mocage french research chemistry model
- NYEAR : year of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read).
- NMONTH : month of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read).
- NDAY : day of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read).
- XTIME : time from midnight of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read). (seconds).
- LWAT_SBL : activates surface boundary multi layer scheme over inland water.

9.5 Lake scheme "FLAKE"

Namelist NAM_PREP_FLAKE

This namelist information is used to initialize the "FLAKE" sea scheme temperature.

Fortran name	Fortran type	values	default value
XTS_WATER_UNIF	real		none
XUNIF_T_SNOW	real		273.15
XUNIF_T_ICE	real		min(273.15,XTS_WATER)
XUNIF_T_MNW	real		depends on XTS_WATER
XUNIF_T_BOT	real		depends on XTS_WATER
XUNIF_T_B1	real		depends on XTS_WATER
XUNIF_H_SNOW	real		0.
XUNIF_H_ICE	real		0. or 0.01 if XTS_WATER < 273.15
XUNIF_H_ML	real		3.
XUNIF_H_B1	real		1.
CFILE_FLAKE	string of 28 characters	'MESONH', 'GRIB '	CFILE in NAM_PREP_SURF_ATM
CTYPE	string of 6 characters		CFILETYPE in NAM_PREP_SURF_ATM
NYEAR	integer		none
NMONTH	integer		none
NDAY	integer		none
XTIME	real		none
LFLK_SBL	logical		F

- XTS_WATER_UNIF : uniform prescribed value of water surface temperature supposed at an **altitude of 0m** (mean sea level altitude). The temperature is then modified for each point depending on its altitude, following a uniform vertical gradient of -6.5 Kkm^{-1} . This prescribed value, if defined, has priority on the use of CFILE_FLAKE data.
- XUNIF_T_SNOW : surface temperature of snow (K)
- XUNIF_T_ICE : surface temperature at the ice-atmosphere or at the ice-snow interface (K)
- XUNIF_T_MNW : mean water column temperature (K)
- XUNIF_T_BOT : water temperature at the bottom of the lake (K)
- XUNIF_T_B1 : temperature at the bottom of the upper layer of sediments (K)
- XUNIF_H_SNOW : snow layer thickness (m)
- XUNIF_H_ICE : ice layer thickness (m)
- XUNIF_H_B1 : thickness of the upper level of the active sediments (m)
- CFILE_FLAKE : name of the file used to define the surface temperature. **The use of a file or prescribed value XTS_WATER_UNIF has priority on the data in CFILE_FLAKE file.**
- CTYPE : type of the CFILE_FLAKE file, if the latter is provided. CTYPE must then be given. The following values are currently usable :
 - "MESONH" : the file type is a MESONH file.
 - "GRIB" : the file type is a GRIB file, coming from any of these models :
 1. "ECMWF" : european center forecast model
 2. "ARPEGE" : Arpege french forecast model
 3. "ALADIN" : Aladin french forecast local model
 4. "MOCAGE" : Mocage french research chemistry model
- NYEAR : year of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read).
- NMONTH : month of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read).
- NDAY : day of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read).
- XTIME : time from midnight of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read). (seconds).
- LFLK_SBL : activates surface boundary multi layer scheme over inland water.

9.6 Vegetation scheme "ISBA "

Namelist NAM_PREP_ISBA

This namelist information is used to initialize the "ISBA " vegetation scheme variables : soil temperature profile, soil water and ice profiles, water intercepted by leaves, snow.

Fortran name	Fortran type	values	default value
XHUG_SURF	real	'MESONH', 'GRIB '	none
XHUG_ROOT	real		none
XHUG_DEEP	real		none
CFILE_WG	string of 28 characters		CFILE_ISBA in this namelist
CTYPE_WG	string of 6 characters		CTYPE in this namelist
XTG_SURF	real	'MESONH', 'GRIB '	none
XTG_ROOT	real		none
XTG_DEEP	real		none
CFILE_TG	string of 28 characters		CFILE_ISBA in this namelist
CTYPE_TG	string of 6 characters		CTYPE in this namelist
CFILE_ISBA	string of 28 characters	'MESONH', 'GRIB '	CFILE in NAM_PREP_SURF_ATM
CTYPE	string of 6 characters		CFILETYPE in NAM_PREP_SURF_ATM
NYEAR	integer		none
NMONTH	integer		none
NDAY	integer		none
XTIME	real		none
LISBA_CANOPY	logical		F

- XHUG_SURF : uniform prescribed value of soil water index (SWI) for the surface soil layer. This prescribed value, if defined, has priority on the use of CFILE_WG and CFILE_ISBA data.
- XHUG_ROOT : uniform prescribed value of soil water index (SWI) for the root zone soil layer(s). This prescribed value, if defined, has priority on the use of CFILE_WG and CFILE_ISBA data.
- XHUG_DEEP : uniform prescribed value of soil water index (SWI) for the deep soil layer(s). This prescribed value, if defined, has priority on the use of CFILE_WG and CFILE_ISBA data.
- CFILE_WG : name of the file used to define the soil water profiles. **The use of a file or prescribed value of XHUG_SURF, XHUG_ROOT and XHUG_DEEP has priority on the data in CFILE_WG file.**
- CTYPE_WG : type of the CFILE_WG file, if the latter is provided. CTYPE_WG must then be given. The following values are currently usable :
 - "MESONH" : the file type is a MESONH file.
 - "GRIB " : the file type is a GRIB file, coming from any of these models :
 1. "ECMWF" : european center forecast model
 2. "ARPEGE" : Arpege french forecast model
 3. "ALADIN" : Aladin french forecast local model
 4. "MOCAGE" : Mocage french research chemistry model
- XTG_SURF : uniform prescribed value of temperature for the surface soil layer, supposed at an **altitude of 0m** (mean sea level altitude). The temperature is then modified for each point depending on its altitude, following a uniform vertical gradient of -6.5 Kkm^{-1} . This prescribed value, if defined, has priority on the use of CFILE_TG and CFILE_ISBA data.
- XTG_ROOT : uniform prescribed value of temperature for the root zone soil layer(s), supposed at an **altitude of 0m** (mean sea level altitude). The temperature is then modified for each point depending on its altitude, following a uniform vertical gradient of -6.5 Kkm^{-1} . This prescribed value, if defined, has priority on the use of CFILE_TG and CFILE_ISBA data.
- XTG_DEEP : uniform prescribed value of temperature for the deep soil layer(s), supposed at an **altitude of 0m** (mean sea level altitude). The temperature is then modified for each point depending on its altitude, following a uniform vertical gradient of -6.5 Kkm^{-1} . This prescribed value, if defined, has priority on the use of CFILE_TG and CFILE_ISBA data.
- CFILE_TG : name of the file used to define the soil temperature profile. **The use of a file or prescribed value of XTG_SURF, XTG_ROOT and XTG_DEEP has priority on the data in CFILE_TG file.**
- CTYPE_TG : type of the CFILE_TG file, if the latter is provided. CTYPE_TG must then be given. The following values are currently usable :
 - "MESONH" : the file type is a MESONH file.
 - "GRIB " : the file type is a GRIB file, coming from any of these models :
 1. "ECMWF" : european center forecast model
 2. "ARPEGE" : Arpege french forecast model
 3. "ALADIN" : Aladin french forecast local model
 4. "MOCAGE" : Mocage french research chemistry model
- CFILE_ISBA : name of the file used to define any ISBA variable. **The use of a file or prescribed value XHUG_SURF, XHUG_ROOT, XHUG_DEEP, XTG_SURF, XTG_ROOT, XTG_DEEP, CFILE_WG and CFILE_TG has priority on the data in CFILE_ISBA file.**
- CTYPE : type of the CFILE_ISBA file, if the latter is provided. CTYPE must then be given. The following values are currently usable :
 - "MESONH" : the file type is a MESONH file.
 - "GRIB " : the file type is a GRIB file, coming from any of these models :
 1. "ECMWF" : european center forecast model
 2. "ARPEGE" : Arpege french forecast model
 3. "ALADIN" : Aladin french forecast local model
 4. "MOCAGE" : Mocage french research chemistry model
- NYEAR : year of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read).
- NMONTH : month of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read).
- NDAY : day of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read).
- XTIME : time from midnight of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read). (seconds).
- LISBA_CANOPY : activates surface boundary multi layer scheme over vegetation.

Namelist NAM_PREP_ISBA_SNOW

This namelist defines the type of snow scheme used in ISBA scheme.

Fortran name	Fortran type	values	default value
CSNOW	string of 3 characters	'D95', '3-L', 'EBA'	'D95'

- CSNOW : type of snow scheme. Possible snow schemes are :
 1. 'D95' : Douville et al (1995) snow scheme.
 2. '3-L' : Boone and Etchevers (2000) three layers snow scheme.
 3. 'EBA' : Bogatchev and Bazile (2005), Arpege operational snow scheme.

9.7 Town scheme "TEB "

Namelist NAM_PREP_TEB

This namelist information is used to initialize the "TEB " urban scheme variables : road, roof and wall temperature profiles, water intercepted by roofs and roads, snow, building internal temperature.

Fortran name	Fortran type	values	default value
XWS_ROAD	real		none
XWS_ROOF	real		none
CFILE_WS	string of 28 characters		CFILE_TEB in this namelist
CTYPE_WS	string of 6 characters	'MESONH', 'GRIB '	CTYPE in this namelist
XTS_ROAD	real		none
XTS_ROOF	real		none
XTS_WALL	real		none
XTI_BLD	real		none
XTI_ROAD	real		none
CFILE_TS	string of 28 characters		CFILE_TEB in this namelist
CTYPE_TS	string of 6 characters	'MESONH', 'GRIB '	CTYPE in this namelist
CFILE_TEB	string of 28 characters		CFILE in NAM_PREP_SURF_ATM
CTYPE	string of 6 characters	'MESONH', 'GRIB '	CFILETYPE in NAM_PREP_SURF_ATM
NYEAR	integer		none
NMONTH	integer		none
NDAY	integer		none
XTIME	real		none
LTEB_CANOPY	logical		F

- XWS_ROAD : uniform prescribed value of soil water interception for the road reservoir. This prescribed value, if defined, has priority on the use of CFILE_WS and CFILE_TEB data.
- XWS_ROOF : uniform prescribed value of soil water interception for the roof reservoir. This prescribed value, if defined, has priority on the use of CFILE_WS and CFILE_TEB data.
- CFILE_WS : name of the file used to define the soil water reservoirs. **The use of a file or prescribed value of XWS_ROAD and XWS_ROOF has priority on the data in CFILE_WS file.**
- CTYPE_WS : type of the CFILE_WS file, if the latter is provided. CTYPE_WS must then be given. The following values are currently usable :
 - "MESONH" : the file type is a MESONH file.
 - "GRIB" : the file type is a GRIB file, coming from any of these models :
 1. "ECMWF" : european center forecast model
 2. "ARPEGE" : Arpege french forecast model
 3. "ALADIN" : Aladin french forecast local model
 4. "MOCAGE" : Mocage french research chemistry model
- XTS_ROAD : uniform prescribed value of surface temperature for road, supposed at an altitude of 0m (mean sea level altitude). The temperature is then modified for each point depending on its altitude, following a uniform vertical gradient of -6.5 Kkm^{-1} . This prescribed value, if defined, has priority on the use of CFILE_TS and CFILE_TEB data.
- XTS_ROOF : uniform prescribed value of surface temperature for roof, supposed at an altitude of 0m (mean sea level altitude). The temperature is then modified for each point depending on its altitude, following a uniform vertical gradient of -6.5 Kkm^{-1} . This prescribed value, if defined, has priority on the use of CFILE_TS and CFILE_TEB data.
- XTS_WALL : uniform prescribed value of surface temperature for wall, supposed at an altitude of 0m (mean sea level altitude). The temperature is then modified for each point depending on its altitude, following a uniform vertical gradient of -6.5 Kkm^{-1} . This prescribed value, if defined, has priority on the use of CFILE_TS and CFILE_TEB data.
- XTI_BLD : uniform prescribed value of internal building temperature. This temperature is not dependent on altitude. This prescribed value, if defined, has priority on the use of CFILE_TS and CFILE_TEB data.
- XTI_ROAD : uniform prescribed value of deep road temperature, supposed at an altitude of 0m (mean sea level altitude). The temperature is then modified for each point depending on its altitude, following a uniform vertical gradient of -6.5 Kkm^{-1} . This prescribed value, if defined, has priority on the use of CFILE_TS and CFILE_TEB data.
- CFILE_TS : name of the file used to define the soil temperature profile. **The use of a file or prescribed value of XTS_ROAD, XTS_ROOF, XTS_WALL, XTI_BLD or XTI_ROAD has priority on the data in CFILE_TS file.**
- CTYPE_TS : type of the CFILE_TS file, if the latter is provided. CTYPE_TS must then be given. The following values are currently usable :
 - "MESONH" : the file type is a MESONH file.
 - "GRIB" : the file type is a GRIB file, coming from any of these models :
 1. "ECMWF" : european center forecast model
 2. "ARPEGE" : Arpege french forecast model
 3. "ALADIN" : Aladin french forecast local model
 4. "MOCAGE" : Mocage french research chemistry model
- CFILE_TEB : name of the file used to define any TEB variable. **The use of a file or prescribed value XWS_ROAD, XWS_ROOF, XTS_ROAD, XTS_ROOF, XTS_WALL, XTI_BLD, XTI_ROAD, CFILE_WS or CFILE_TS has priority on the data in CFILE_TEB file.**
- CTYPE : type of the CFILE_TEB file, if the latter is provided. CTYPE must then be given. The following values are currently usable :
 - "MESONH" : the file type is a MESONH file.
 - "GRIB" : the file type is a GRIB file, coming from any of these models :
 1. "ECMWF" : european center forecast model
 2. "ARPEGE" : Arpege french forecast model
 3. "ALADIN" : Aladin french forecast local model
 4. "MOCAGE" : Mocage french research chemistry model
- NYEAR : year of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read).
- NMONTH : month of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read).
- NDAY : day of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read).
- XTIME : time from midnight of surface UTC time. It is used only if no atmospheric file or no surface file is given (in those the date can be read). (seconds).
- LTEB_CANOPY : activates surface boundary multi layer scheme over town.

Chapitre 10

How to run the externalized surface physical schemes

Here are described the options available during the run of the several surface schemes.

10.1 "SURF_ATM" general options available over all tiles

Namelist NAM_SURF_ATM

Fortran name	Fortran type	values	default value
XCISMIN	real		6.7E-5
XVMODMIN	real		0.
LALDTHRES	logical		F
LALDZ0H	logical		F
LDRAG_COEF_ARP	logical		F
XEDB	real		5.
XEDC	real		5.
XEDD	real		5.
XEDK	real		1.
XUSURIC	real		1.
XUSURID	real		0.035
XUSURICL	real		4.
XVCHRNK	real		0.015
XVZ0CM	real		0.0

- LALDTHRES : flag to set a minimum wind and shear like done in Aladin model.
- XCISMIN : minimum wind shear to compute turbulent exchange coefficient (used only if LALDTHRES)
- XVMODMIN : minimum wind speed to compute turbulent exchange coefficient (used only if LALDTHRES)
- LALDZ0H : to take into account orography in heat roughness length
- XEDB, XEDC, XEDD, XEDK : coefficients used in Richardson critical numbers computation
- XUSURIC, XUSURID, XUSURICL : Richardson critical numbers
- XVCHRNK, XVZ0CM : Charnock's constant and minimal neutral roughness length over sea (formulation of roughness length over sea)

10.2 "SEAFLX" sea scheme options

Namelist NAM_SEAFLUXn

Fortran name	Fortran type	values	default value
CSEA_FLUX	string of 6 characters	'DIRECT', 'ITERAT', 'COARE3', 'ECUME '	'ECUME '
CSEA_ALB	string of 4 characters	'UNIF', 'TA96'	'TA96'
LPWG	logical		F
LPRECIP	logical		F
LPWEBB	logical		F

- CSEA_FLUX : type of flux computation physics. The following option is currently available :
 - "DIRECT" : direct Charnock computation. No effect of convection in the the boundary layer on the fluxes formulae.
 - "ITERAT" : iterative method proposed by Fairall et al (1996) from TOGA-COARE experiment, amended by Mondon and Redelsperger (1998) to take into account effect of atmospheric convection on fluxes.
 - "COARE3" : iterative method proposed by Fairall et al (1996) from TOGA-COARE experiment, amended by cnrm/memo to take into account effect of atmospheric convection, precipitation and gustiness on fluxes.
 - "ECUME " : iterative method proposed by Fairall et al (1996) from TOGA-COARE experiment, amended by cnrm/memo to take into account effect of atmospheric convection, precipitation and gustiness on fluxes : improvement of surface exchange coefficients representation.
 - LPWG : correction of fluxes due to gustiness
 - LPRECIP : correction of fluxes due to precipitation
 - LPWEBB : correction of fluxes due to convection (Webb effect)
- CSEA_ALB : type of albedo formula. The following options are currently available :
 - "UNIF" : a uniform value of 0.135 is used for water albedo
 - "TA96" : Taylor et al (1996) formula for water direct albedo, depending on solar zenith angle θ : $\alpha_{dir} = 0.037 / \left(1.1 \cos(\theta)^{1.4} + 0.15 \right)$

10.3 "FLAKE" lake scheme options

Namelist NAM_FLAKEn

Fortran name	Fortran type	values	default value
LSEDIMENTS	logical		T
CFLAKE_SNOW	string of 6 characters	'NON ', 'FLAKE ', 'ISBAES'	'NON '
CFLAKE_FLUX	string of 6 characters	'FLAKE ', 'ISBA '	'FLAKE '

- CFLAKE_SNOW : snow scheme to be used. For the time being only option 'FLAKE ' is active
- CFLAKE_FLUX : scheme to be used to compute surface fluxes of moment, energy and water vapor. For the time being only option 'FLAKE ' is active
- LSEDIMENTS : to use the bottom sediments scheme of Flake (default)

10.4 "ISBA" vegetation scheme options

Namelist NAM_SGH_ISBA

Fortran name	Fortran type	values	default value
CRUNOFF	string of 4 characters	'WSAT', 'DT92', 'SGH'	'WSAT'
CTOPREG	string of 4 characters	'DEF', 'NON'	'DEF'
CKSAT	string of 4 characters	'DEF', 'SGH'	'DEF'
CRAIN	string of 3 characters	'DEF', 'SGH'	'DEF'
CHORT	string of 4 characters	'DEF', 'SGH'	'DEF'

- CRUNOFF : type of subgrid runoff. The following options are currently available :
 - "WSAT" : runoff occurs only when saturation is reached
 - "DT92" : Dumenill and Todini (1992) subgrid runoff formula
 - "SGH" : Decharme et al. (2006) Topmodel like subgrid runoff
- CTOPREG : kind of regression. Option activated only if CRUNOFF = 'SGH'. The following options are currently available :
 - "DEF" : Wolock and MacCabe regression between topographic indices computed at 1km and 100m resolution (recommended)
 - "NON" : no regression
- CKSAT : Activates the exponential profile for Ksat. The following options are currently available :
 - "DEF" : homogeneous profile
 - "SGH" : exponential decreasing profile with depth (due to compaction of soil)
- CRAIN : Activates the spatial distribution of rainfall intensity. The following options are currently available :
 - "DEF" : homogeneous distribution
 - "SGH" : exponential distribution which depends on the fraction of the mesh where it rains. This fraction depends on the mesh resolution and the intensity of hourly precipitation. (If the horizontal mesh is lower than 10km then the fraction equals 1).
- CHORT : Activates the Horton runoff due to water infiltration excess. The following options are currently available :
 - "DEF" : no Horton runoff
 - "SGH" : Horton runoff computed

Namelist NAM_ISBA

Fortran name	Fortran type	values	default value
CCIDRY	string of 4 characters	'DEF', 'GB93'	'DEF'
CSCOND	string of 4 characters	'NP89', 'PL98'	'NP89'
CSOILFRZ	string of 3 characters	'DEF', 'LWT'	'DEF'
CDIFSFCOND	string of 4 characters	'DEF', 'MLCH'	'DEF'
CSNOWRES	string of 3 characters	'DEF', 'RIL'	'DEF'
CALBEDO	string of 4 characters	'MEAN', 'DRY', 'WET', 'EVOL'	'DRY'
CROUGH	string of 4 characters	'Z01D', 'Z04D'	'Z04D'
CCPSURF	string of 3 characters	'DRY', 'HUM'	'DRY'
XTSTEP	real		none
XCGMAX	real		2.E-5

- CCIDRY : type of C1 formulation for dry soils. The following options are currently available :
 - "DEF" : Giard-Bazile formulation
 - "GB93" : Giordani 1993, Braud 1993
- CSCOND : type of thermal conductivity. The following options are currently available :
 - "NP89" : Noilhan and Planton (1989) formula
 - "PL98" : Peters-Lidard et al. (1998) formula
- CSOILFRZ : type of soil freezing-physics option. The following options are currently available :
 - "DEF" : Boone et al. 2000 ; Giard and Bazile 2000
 - "LWT" : Phase changes as above, but relation between unfrozen water and temperature considered
- CDIFSFCOND : type of Mulch effects. The following options are currently available :
 - "DEF" : no mulch effect
 - "MLCH" : include the insulating effect of leaf litter/mulch on the surf. thermal cond.
- CSNOWRES : type of turbulent exchanges over snow. The following options are currently available :
 - "DEF" : Louis
 - "RIL" : Maximum Richardson number limit for stable conditions ISBA-SNOW3L turbulent exchange option
- CALBEDO : type of bare soil albedo. The following options are currently available :
 - "DRY" : dry bare soil albedo
 - "WET" : wet bare soil albedo
 - "MEAN" : albedo for bare soil half wet, half dry
 - "EVOL" : albedo of bare soil evolving with soil humidity
- CROUGH : type of orographic roughness length. The following options are currently available :
 - "Z01D" : orographic roughness length does not depend on wind direction
 - "Z04D" : orographic roughness length depends on wind direction
- CCPSURF : type of specific heat at surface. The following options are currently available :
 - "DRY" : specific heat does not depend on humidity at surface
 - "HUM" : specific heat depends on humidity at surface.
- XTSTEP : time step for ISBA. Default is to use the time-step given by the atmospheric coupling (seconds).
- XCGMAX : maximum value for soil heat capacity.

Namelist NAM_SURF_DST

Fortran name	Fortran type	values	default value
CEMISPARAM	string of 5 characters	'Dal87'	'Dal87'
CVERMOD	string of 6 characters	'CMDVER'	'CMDVER'

- "CEMISPARAM" : parameterization type

Chapitre 11

How to run the externalized surface chemical schemes

Here are described the options available during the run of the several schemes for emission and deposition of chemical species. Note that all the schemes for deposition and emission of chemical species do activate only if chemical species are present (i.e. if the coupling between atmosphere and surface include the chemical species concentrations and fluxes).

11.1 Chemical settings control

Namelist NAM_CH_CONTROLn

Fortran name	Fortran type	values	default value
CCHEM_SURF_FILE	string of 28 characters		' '

- CCHEM_SURF_FILE : name of general (chemical) purpose ASCII input file.

11.2 Chemical anthropogenic emissions

Namelist NAM_CH_SURFn

Fortran name	Fortran type	values	default value
LCH_SURF_EMIS	logical		.FALSE.

- LCH_SURF_EMIS : flag to use anthropogenic emissions or not.

11.3 Chemical deposition over ocean

Namelist NAM_CH_SEAFLUXn

Fortran name	Fortran type	values	default value
CCH_DRY_DEP	string of 6 characters	'NONE ', 'WES89 '	'WES89 '

- CCH_DRY_DEP : type of deposition scheme.
- "NONE " : no chemical deposition scheme.
- "WES89 " : Wesley (1989) deposition scheme.

11.4 Chemical deposition over lakes

Namelist NAM_CH_WATFLUXn

Fortran name	Fortran type	values	default value
CCH_DRY_DEP	string of 6 characters	'NONE ', 'WES89 '	'WES89 '

- CCH_DRY_DEP : type of deposition scheme.
- "NONE " : no chemical deposition scheme.
- "WES89 " : Wesley (1989) deposition scheme.

11.5 Chemical deposition over towns

Namelist NAM_CH_TEBn

Fortran name	Fortran type	values	default value
CCH_DRY_DEP	string of 6 characters	'NONE ', 'WES89 '	'WES89 '

- CCH_DRY_DEP : type of deposition scheme.
- "NONE " : no chemical deposition scheme.
- "WES89 " : Wesley (1989) deposition scheme.

11.6 Chemical deposition and biogenic emissions over vegetation

Namelist NAM_CH_ISBAn

Fortran name	Fortran type	values	default value
CCH_DRY_DEP	string of 6 characters	'NONE ', 'WES89 '	'WES89 '
LCH_BIO_FLUX	logical		.FALSE.

- CCH_DRY_DEP : type of deposition scheme.
- "NONE " : no chemical deposition scheme.
- "WES89 " : Wesley (1989) deposition scheme.
- LCH_BIO_FLUX : flag to activate the biogenic emissions.

Chapitre 12

Externalized surface diagnostics

The diagnostics for the surface require the call to the complete physics of the surface. Therefore, they can be computed either during the run of the schemes (in order to have for example continuous time series of these diagnostics), or can be computed at a given instant only, if atmospheric forcing is given at this instant for the surface scheme to do one time step. The cumulated diagnostics are of course significant only when computed during a run.

12.1 Diagnostics relative to the general surface monitor

Namelist NAM_DIAG_SURF_ATMn

Fortran name	Fortran type	default value
LFRAC	logical	.FALSE.

- LFRAC : flag to save in the output file the sea, inland water, natural covers and town fractions.

12.2 Diagnostics relative to the general surface monitor and to each surface scheme

Namelist NAM_DIAG_SURFn

Fortran name	Fortran type	values	default value
N2M	integer	0, 1, 2	0
LSURF_BUDGET	logical		.FALSE.
LRAD_BUDGET	logical		.FALSE.
LCOEF	logical		.FALSE.
LSURF_VARS	logical		.FALSE.

- N2M : flag to compute surface boundary layer characteristics :
 - N2M=1 : computes temperature at 2 m, specific humidity at 2 m, relative humidity, zonal and meridian wind at 10 m, and Richardson number. 2m and 10m quantities are calculated extrapolating atmospheric forcing variables with Paulson laws using surface heat, water and momentum fluxes.
 - N2M=2 : computes temperature at 2 m, specific humidity at 2 m, relative humidity, zonal and meridian wind at 10 m, and Richardson number. 2m and 10m quantities are calculated interpolating between atmospheric forcing variables and surface temperature and humidity.
- LSURF_BUDGET : flag to save in the output file the terms of the surface energy balance (net radiation, sensible heat flux, latent heat flux, ground flux), for each scheme (on the four separate tiles), on each patch of the vegetation scheme if existing, and aggregated for the whole surface. The diagnosed fields are (* stands for the scheme considered (*=nothing : field aggregated on the whole surface ; *=name of a scheme : field for this scheme) :
 - RN_* : net radiation
 - H_* : turbulent sensible heat flux
 - LE_* : turbulent latent heat flux
 - GFLUX_* : ground or storage heat flux
 - SWD_* : downward short wave radiation
 - SWU_* : upward short wave radiation
 - SWBD_* : downward short wave radiation for each spectral band
 - SWBU_* : upward short wave radiation for each spectral band
 - LWD_* : downward long wave radiation
 - LWU_* : upward long wave radiation
 - FMU_* : zonal wind stress
 - FMV_* : meridian wind stress
 - Z0EFF : effective roughness length (vegetation and orography)
- LRAD_BUDGET : flag used to write downward and upward shortwave radiation per spectral band into output file (they're computed even if LRAD_BUDGET is false)
- LCOEF : flag to save in the output file the transfer coefficients used in the computation of the surface energy fluxes, for each scheme (on the four separate tiles) and aggregated for the whole surface. The diagnosed fields are (* stands for the scheme considered (*=nothing : field aggregated on the whole surface ; *=name of a scheme : field for this scheme) :
 - CD_* : drag coefficient for momentum
 - CH_* : drag coefficient for heat
 - CE_* : drag coefficient for evaporation (differs from CH only over sea)
 - Z0_* : roughness length
 - Z0H_* : thermal roughness length
- LSURF_VARS : flag to save in the output file the surface specific humidity for each scheme (on the four separate tiles), on each patch of the vegetation scheme if existing. The diagnosed fields are (* stands for the scheme considered (*=nothing : field aggregated on the whole surface ; *=name of a scheme : field for this scheme) :
 - QS_* : specific humidity

12.3 Diagnostics relative to the ISBA vegetation scheme

Namelist NAM_DIAG_ISBAn

Fortran name	Fortran type	default value
LPGD	logical	.FALSE.
LSURF_EVAP_BUDGET	logical	.FALSE.
LSURF_MISC_BUDGET	logical	.FALSE.
LSURF_BUDGETC	logical	.FALSE.

- LPGD : flag to save in the output file the physiographic fields of ISBA scheme that are computed from ecoclimap data from the ecosystem fractions.
- LSURF_EVAP_BUDGET : flag to save in the output file the detailed terms of the water vapor fluxes, on each patch of the vegetation scheme if existing, and aggregated for the natural surface. The diagnosed fields are :
- LSURF_MISC_BUDGET : flag to save in the output file miscellaneous fields. The diagnosed fields are :
 - HV : Halstead coefficient
 - SNG : snow fraction over bare ground
 - SNV : snow fraction over vegetation
 - SN : total snow fraction
 - SWI : soil wetness index for each ground layer $(w_g - w_{wilt})/(w_{fc} - w_{wilt})$ where w_g is the volumic water content, w_{fc} is the porosity and w_{wilt} corresponds to the plant wilting point.
 - GPP : Gross primary production
 - RDK : Dark respiration
- LSURF_BUDGETC : flag to save in the output file the time integrated values of all budget terms that have been activated (either with :LSURF_BUDGET flag in NAM_DIAG_SURF_n or LSURF_EVAP_BUDGET in the present namelist).

12.4 Diagnostics relative to the TEB town scheme

Namelist NAM_DIAG_TEBn

Fortran name	Fortran type	default value
LSURF_MISC_BUDGET	logical	.FALSE.

- LSURF_MISC_BUDGET : flag to save in the output file miscellaneous fields. The diagnosed fields are :
 - ZO_TOWN : roughness length for town
 - QF_BLD : domestic heating
 - QF_BLDWFR : domestic heating
 - FLX_BLD : heat flux from bld
 - TL_BLD_EQ : internal temperature without heating
 - TL_BLDWFR : internal temperature without heating
 - QF_TOWN : total anthropogenic heat
 - DQS_TOWN : storage inside building
 - H_WALL : wall sensible heat flux
 - H_ROOF : roof sensible heat flux
 - H_ROAD : road sensible heat flux
 - RN_WALL : net radiation at wall
 - RN_ROOF : net radiation at roof
 - RN_ROAD : net radiation at road
 - GFLUX_WALL : net wall conduction flux
 - GFLUX_ROOF : net roof conduction flux
 - GFLUX_ROAD : net road conduction flux
 - LE_ROOF : roof latent heat flux
 - LE_ROAD : road latent heat flux

Chapitre 13

Externalized surface model output fields

Model output fields depend on the tile and on the configuration of run.

13.1 Prognostic model output fields

13.1.1 ISBA

The definition of the representation of soil, vegetation, snow and surface boundary layer components is done during PGD and PREP. The description of soil is done with parameter CISBA (2 layers, 3 layers or more if diffusion treatment selected) from NAM_ISBA. The representation of vegetation is done with parameters NPATCH (number of patches over vegetation) and CPHOTO (type of photosynthesis). During PREP, the snow scheme is chosen by parameter CSNOW in NAM_PREP_ISBA and if the surface boundary layer (SBL) scheme is activated then LISBA_CANOPY key has to be set to T.

[illegible]

variable	dimension	unit	description
TG1	[2D]	[K]	surface temperature
TG2	[2D]	[K]	deep soil temperature
TG3	[2D]	[K]	third layer temperature
WG1	[2D]	[m3/m3]	surface liquid volumetric water content
WG2	[2D]	[m3/m3]	root liquid volumetric water content
WG3	[2D]	[m3/m3]	deep liquid volumetric water content
WGI1	[2D]	[m3/m3]	surface frozen volumetric water content
WGI2	[2D]	[m3/m3]	root frozen volumetric water content
WGI3	[2D]	[m3/m3]	third layer frozen volumetric water content
WR	[2D]	[Kg/m2]	liquid water retained by the foliage
RESA	[2D]	[s/m]	aerodynamical resistance
LAI	[2D]	[m2/m2]	leaf area index
AN	[2D]	[kgCO2/m2/s]	CO2 net assimilation
ANDAY	[2D]	[kgCO2/m2]	daily CO2 net assimilation
ANFM	[2D]	[mg/m2/s]	maximum leaf assimilation
LE_AGS	[2D]	[W/m2]	latent heat flux
RESPBSTR	[2D]	[W/m2]	respiration of above-ground structural biomass
RESPBSTR2	[2D]	[W/m2]	respiration of deep structural biomass
BIOMASSTR2	[2D]	[Kg/m2]	total dry above-ground structural biomass
BIOMASSTR2_LAST	[2D]	[Kg/m2]	total dry below-ground structural biomass of day-1
BIOMASSTR_LAST	[2D]	[Kg/m2]	total dry above-ground structural biomass of day-1
TSNOW_VEG1	[2D]	[K]	1st layer snow temperature
WSNOW_VEG1	[2D]	[Kg/m2]	1st layer snow water equivalent
RSNOW_VEG1	[2D]	[Kg/m3]	1st layer snow density
HSNOW_VEG1	[2D]	[W/m2]	1st layer snow heat content
ASNOW_VEG	[2D]	[-]	snow albedo
WSNOW_VEG2	[2D]	[Kg/m2]	2nd layer snow water equivalent
RSNOW_VEG2	[2D]	[Kg/m3]	2nd layer snow density
HSNOW_VEG2	[2D]	[W/m2]	2nd layer snow heat content
WSNOW_VEG3	[2D]	[Kg/m2]	3rd layer snow water equivalent
RSNOW_VEG3	[2D]	[Kg/m3]	3rd layer snow density
HSNOW_VEG3	[2D]	[W/m2]	3rd layer snow heat content
ISBA_CAN_Un	[1D]	[m/s]	wind in canopy at layer n (n=1 to 6)
ISBA_CAN_Tn	[1D]	[m/s]	temperature in canopy at layer n (n=1 to 6)
ISBA_CAN_Qn	[1D]	[m/s]	humidity in canopy at layer n (n=1 to 6)
ISBA_CAN_En	[1D]	[m/s]	TKE in canopy at layer n (n=1 to 6)
ISBA_CAN_Pn	[1D]	[m/s]	pressure in canopy at layer n (n=1 to 6)

13.1.2 SEAFLUX

	SEAFLUX	SBL
SST	•	
Z0SEA	•	
SEA_SBL_U		•
SEA_SBL_T		•
SEA_SBL_Q		•
SEA_SBL_E		•
SEA_SBL_P		•

13.1.3 TEB

name	Standard TEB		CANOPY
	layer		
T_ROOF	1	•	
	2	•	
	3	•	
WS_ROOF		•	
T_ROAD	1	•	
	2	•	
	3	•	
WS_ROAD		•	
T_WALL	1	•	
	2	•	
	3	•	
TL_BLD		•	
TL_ROAD		•	
WSNOW_ROOF	1	•	
RSNOW_ROOF	1	•	
TSNOW_ROOF	1	•	
ASNOW_ROOF		•	
WSNOW_ROAD	1	•	
RSNOW_ROAD	1	•	
TSNOW_ROAD	1	•	
ASNOW_ROAD		•	
T_CANYON		•	
Q_CANYON		•	
TEB_CAN_U			•
TEB_CAN_T			•
TEB_CAN_Q			•
TEB_CAN_E			•
TEB_CAN_P			•

13.1.4 WATFLUX

	WATERFLUX	SBL
TS_WATER	•	
Z0WATER	•	
WAT_SBL_U		•
WAT_SBL_T		•
WAT_SBL_Q		•
WAT_SBL_E		•
WAT_SBL_P		•

13.1.5 FLAKE

	FLAKE	SBL
TS_WATER	•	
T_SNOW	•	
T_ICE	•	
T_MNW	•	
T_WML	•	
T_BOT	•	
T_BI	•	
CT	•	
H_SNOW	•	
H_ICE	•	
H_ML	•	
H_BI	•	
WAT_SBL_U		•
WAT_SBL_T		•
WAT_SBL_Q		•
WAT_SBL_E		•
WAT_SBL_P		•

Annexe A

Example of namelist

A.1 surface parameters defined from ecoclimap

This is the classical way how to use ecoclimap and other databases (orography, sand and clay). The initialization of prognostic variables are done by reading a grib file (Arpege file in the example).

```
&NAM_PGDFILE          CPGDFILE = 'PGDFILE'
/
&NAM_FRAC              LECOCLIMAP = T
/
&NAM_PGD_GRID          CGRID = 'LONLAT REG'
/
&NAM_CONF_PROJ         XLATO = -90.          ,
                        XLONO = 0.            ,
                        XRPK  = -1.            ,
                        XBETA = 0.             
/
&NAM_CONF_PROJ_GRID    NIMAX  = 200          ,
                        NJMAX  = 200          ,
                        XLATCEN = -90.         ,
                        XLONCEN = 0.           ,
                        XDX     = 25000.       ,
                        XDY     = 25000.      
/
&NAM_LONLAT_REG         XLONMIN = -4.5        ,
                        XLONMAX = -4.          ,
                        XLATMIN = 48.          ,
                        XLATMAX = 48.5         ,
                        NLON    = 6            ,
                        NLAT    = 7            
/
&NAM_PGD_SCHEMES        CNATURE = 'ISBA'      ,
                        CSEA    = 'SEAFLX'    ,
                        CTOWN   = 'TEB'       ,
                        CWATER  = 'WATFLX'    
/
&NAM_COVER              YCOVER  = 'ecoclimats_v2' ,
                        YFILETYPE = 'DIRECT'
/
&NAM_ZS                 YZS      = 'gtopo30'    ,
                        YFILETYPE = 'DIRECT'
/
&NAM_ISBA               YCLAY    = 'clay_fao'    ,
                        YCLAYFILETYPE = 'DIRECT' ,
                        YSAND     = 'sand_fao'    ,
                        YSANDFILETYPE = 'DIRECT' ,
                        XUNIF_RUNOFFB = 0.5        ,
                        CISBA     = '3-L'         ,
                        CPHOTO    = 'NON'         ,
                        NPATCH    = 1              ,
                        NGROUND_LAYER = 3
```

/

```
&NAM_PREPFILE      CPREPFILE = 'PREP'
/
&NAM_PREP_SURF_ATM  CFILE       = 'arpifs.AN.20030101.00' ,
                     CFILETYPE    = 'GRIB  '
/
&NAM_PREP_TEB       CFILE_TEB    = 'arpifs.AN.20030101.00' ,
                     CTYPE        = 'GRIB  '
/
&NAM_PREP_SEAFLUX   CFILE_SEAFLX = 'arpifs.AN.20030101.00' ,
                     CTYPE        = 'GRIB  '
/
&NAM_PREP_WATFLUX   CFILE_WATFLX = 'arpifs.AN.20030101.00' ,
                     CTYPE        = 'GRIB  '
/
&NAM_PREP_ISBA      CFILE_ISBA   = 'arpifs.AN.20030101.00' ,
                     CTYPE        = 'GRIB  '
/
&NAM_PREP_ISBA_SNOW CSNOW       = '3-L'
```

```
&NAM_IO_OFFLINE     YPROGRAM    = 'ASCII ' ,
                     YCOUPLING   = 'E'
/
&NAM_DIAG_SURF_n    LSURF_BUDGET = T      ,
                     N2M         = 1      ,
                     LCOEF       = T
/
&NAM_DIAG_SURF_ATM_n LFRAC       = T
/
&NAM_DIAG_ISBA_n    LPGD          = T ,
                     LSURF_EVAP_BUDGET = T ,
                     LSURF_MISC_BUDGET = T ,
                     LSURF_BUDGETC    = F
/
&NAM_DIAG_TEB_n     LSURF_MISC_BUDGET = T
/
&NAM_SGH_ISBA_n     CRUNOFF       = "WSAT"
/
&NAM_ISBA_n          CROUGH        = "Z04D" ,
                     CSCOND        = "NP89" ,
                     CALBEDO       = "DRY" ,
                     CC1DRY        = 'DEF' ,
                     CSOILFRZ      = 'DEF' ,
                     CDIFSFCND     = 'DEF' ,
                     CCPSURF       = 'DRY' ,
                     CSNOWRES      = 'DEF'
/
&NAM_CH_ISBA_n      CCH_DRY_DEP   = "WES89 "
/
&NAM_SEAFLUX_n      CSEA_ALB      = "TA96" ,
                     CSEA_FLUX     = "COARE2" ,
                     LPWG          = T      ,
                     LPRECIP       = T      ,
                     LPWEBB        = T
/
&NAM_CH_SEAFLUX_n   CCH_DRY_DEP   = "WES89 "
/
&NAM_CH_WATFLUX_n   CCH_DRY_DEP   = "WES89 "
/
&NAM_CH_TEB_n       CCH_DRY_DEP   = "WES89 "
```

A.2 user defined surface parameters

Ecoclimap is not used (LECOCLIMAP = F). Information is not read from databases but the user defines his own surface parameters. Uniform field is used in this 1D case.

A.2.1 Uniform values prescribed : 1d example without patches

```
&NAM_DATA_ISBA      NTIME = 12 ,
                     XUNIF_VEGTYPE(1)      = 0.,
                     XUNIF_VEGTYPE(2)      = 0.,
                     XUNIF_VEGTYPE(3)      = 0.,
                     XUNIF_VEGTYPE(4)      = 0.,
                     XUNIF_VEGTYPE(5)      = 1.,
                     XUNIF_VEGTYPE(6)      = 0.,
                     XUNIF_VEGTYPE(7)      = 0.,
                     XUNIF_VEGTYPE(8)      = 0.,
                     XUNIF_VEGTYPE(9)      = 0.,
                     XUNIF_VEGTYPE(10)     = 0.,
                     XUNIF_VEGTYPE(11)     = 0.,
                     XUNIF_VEGTYPE(12)     = 0.,
                     XUNIF_VEG(1,1)        = 1.,
                     XUNIF_VEG(1,2)        = 1.,
                     XUNIF_VEG(1,3)        = 1.,
                     XUNIF_VEG(1,4)        = 1.,
                     XUNIF_VEG(1,5)        = 1.,
                     XUNIF_VEG(1,6)        = 1.,
                     XUNIF_VEG(1,7)        = 1.,
                     XUNIF_VEG(1,8)        = 1.,
                     XUNIF_VEG(1,9)        = 1.,
                     XUNIF_VEG(1,10)       = 1.,
                     XUNIF_VEG(1,11)       = 1.,
                     XUNIF_VEG(1,12)       = 1.,
                     XUNIF_LAI(1,1)        = 0.789,
                     XUNIF_LAI(1,2)        = 0.213,
                     XUNIF_LAI(1,3)        = 0.345,
                     XUNIF_LAI(1,4)        = 0.467,
                     XUNIF_LAI(1,5)        = 0.888,
                     XUNIF_LAI(1,6)        = 0.621,
                     XUNIF_LAI(1,7)        = 0.743,
                     XUNIF_LAI(1,8)        = 0.855,
                     XUNIF_LAI(1,9)        = 0.976,
                     XUNIF_LAI(1,10)       = 0.123,
                     XUNIF_LAI(1,11)       = 0.134,
                     XUNIF_LAI(1,12)       = 0.155,
                     XUNIF_ZO(1,1)         = 2.5,
                     XUNIF_ZO(1,2)         = 2.5,
                     XUNIF_ZO(1,3)         = 2.5,
                     XUNIF_ZO(1,4)         = 2.5,
                     XUNIF_ZO(1,5)         = 2.5,
                     XUNIF_ZO(1,6)         = 2.5,
                     XUNIF_ZO(1,7)         = 2.5,
                     XUNIF_ZO(1,8)         = 2.5,
                     XUNIF_ZO(1,9)         = 2.5,
                     XUNIF_ZO(1,10)        = 2.5,
                     XUNIF_ZO(1,11)        = 2.5,
                     XUNIF_ZO(1,12)        = 2.5,
                     XUNIF_EMIS(1,1)       = 0.97,
                     XUNIF_EMIS(1,2)       = 0.97,
                     XUNIF_EMIS(1,3)       = 0.97,
                     XUNIF_EMIS(1,4)       = 0.97,
                     XUNIF_EMIS(1,5)       = 0.97,
                     XUNIF_EMIS(1,6)       = 0.97,
                     XUNIF_EMIS(1,7)       = 0.97,
                     XUNIF_EMIS(1,8)       = 0.97,
                     XUNIF_EMIS(1,9)       = 0.97,
                     XUNIF_EMIS(1,10)      = 0.97,
                     XUNIF_EMIS(1,11)      = 0.97,
                     XUNIF_EMIS(1,12)      = 0.97,
```

```

XUNIF_DG(1,1)      = 0.01,
XUNIF_DG(1,2)      = 0.70,
XUNIF_DG(1,3)      = 1.30,
XUNIF_ROOTFRAC(1,1) = -999.,
XUNIF_ROOTFRAC(1,2) = -999.,
XUNIF_ROOTFRAC(1,3) = -999.,
XUNIF_RSMIN(1)     = 150.,
XUNIF_GAMMA(1)      = 0.04,
XUNIF_WRMX_CF(1)    = 0.1,
XUNIF_RGL(1)        = 30.,
XUNIF_CV(1)         = 0.00001,
XUNIF_ZO_O_ZOH(1)   = 10.,
XUNIF_ALBNIR_VEG(1) = 0.15,
XUNIF_ALBVIS_VEG(1) = 0.05,
XUNIF_ALBUV_VEG(1)  = 0.0425,
XUNIF_ALBNIR_SOIL(1) = 0.15,
XUNIF_ALBVIS_SOIL(1) = 0.05,
XUNIF_ALBUV_SOIL(1) = 0.0425,
XUNIF_GMES(1)       = 0.001,
XUNIF_RE25(1)       = 0.00000015,
XUNIF_BSLAI(1)      = 0.25,
XUNIF_LAIMIN(1)     = 1.0,
XUNIF_SEFOLD(1)     = 31536000.,
XUNIF_GC(1)         = 0.,
XUNIF_DMAX(1)       = 0.1,
XUNIF_F2I(1)        = 0.3,
XUNIF_H_TREE(1)     = 20.,
XUNIF_CE_NITRO(1)   = 4.85,
XUNIF_CF_NITRO(1)   = -0.24,
XUNIF_CNA_NITRO(1)  = 2.8,

```

/
&NAM_DATA_TEB

```

NROOF_LAYER        = 3,
XUNIF_ALB_ROOF     = 0.2,
XUNIF_EMIS_ROOF    = 0.97,
XUNIF_HC_ROOF(1)   = 2110000.,
XUNIF_HC_ROOF(2)   = 2800000.,
XUNIF_HC_ROOF(3)   = 2900000.,
XUNIF_TC_ROOF(1)   = 1.51,
XUNIF_TC_ROOF(2)   = 0.08,
XUNIF_TC_ROOF(3)   = 0.05,
XUNIF_D_ROOF(1)    = 0.05,
XUNIF_D_ROOF(2)    = 0.4,
XUNIF_D_ROOF(3)    = 0.1,
NROAD_LAYER        = 3,
XUNIF_ALB_ROAD     = 0.2,
XUNIF_EMIS_ROAD    = 0.97,
XUNIF_HC_ROAD(1)   = 2110000.,
XUNIF_HC_ROAD(2)   = 2800000.,
XUNIF_HC_ROAD(3)   = 2900000.,
XUNIF_TC_ROAD(1)   = 1.51,
XUNIF_TC_ROAD(2)   = 0.08,
XUNIF_TC_ROAD(3)   = 0.05,
XUNIF_D_ROAD(1)    = 0.05,
XUNIF_D_ROAD(2)    = 0.4,
XUNIF_D_ROAD(3)    = 0.1,
NWALL_LAYER        = 3,
XUNIF_ALB_WALL     = 0.2,
XUNIF_EMIS_WALL    = 0.97,
XUNIF_HC_WALL(1)   = 2110000.,
XUNIF_HC_WALL(2)   = 2800000.,
XUNIF_HC_WALL(3)   = 2900000.,
XUNIF_TC_WALL(1)   = 1.51,
XUNIF_TC_WALL(2)   = 0.08,
XUNIF_TC_WALL(3)   = 0.05,
XUNIF_D_WALL(1)    = 0.05,
XUNIF_D_WALL(2)    = 0.4,
XUNIF_D_WALL(3)    = 0.1,
XUNIF_ZO_TOWN      = 1.,

```

```

XUNIF_BLD          = 0.5,
XUNIF_BLD_HEIGHT = 10.,
XUNIF_WALL_O_HOR  = 0.5,
XUNIF_H_TRAFFIC   = 10.,
XUNIF_LE_TRAFFIC  = 0.,
XUNIF_H_INDUSTRY  = 5.,
XUNIF_LE_INDUSTRY= 0.

/
&NAM_FRAC          LECOCLIMAP = F,
XUNIF_SEA          = 0.,
XUNIF_WATER        = 0.5,
XUNIF_TOWN         = 0.,
XUNIF_NATURE       = 0.5

/
&NAM_PGD_GRID      CGRID = 'LONLAT REG'
/
&NAM_LONLAT_REG    XLONMIN = 0.          ,
XUNIF              XLONMAX = 0.          ,
XUNIF              XLATMIN = 0.          ,
XUNIF              XLATMAX = 0.          ,
NLON               = 1                    ,
NLAT               = 1

/
&NAM_PGD_SCHEMES   CNATURE = 'ISBA'   ,
CSEA               = 'SEAFLX' ,
CTOWN              = 'TEB'    ,
CWATER             = 'WATFLX'

/
&NAM_ZS            XUNIF_ZS = 0.

/
&NAM_ISBA          XUNIF_CLAY = 0.4      ,
XUNIF              XUNIF_SAND = 0.2      ,
XUNIF              XUNIF_RUNOFFB = 0.5    ,
CISBA              = '3-L' ,
CPHOTO             = 'NON' ,
NPATCH             = 1      ,
NGROUND_LAYER      = 3

/

&NAM_PREPFILE      CPREPFILE = 'PREP'
/
&NAM_PREP_SURF_ATM NYEAR = 2004,
NMONTH = 10,
NDAY = 25,
XTIME = 21600.

/
&NAM_PREP_SEAFLUX  XSST_UNIF = 285.,
NYEAR = 2004,
NMONTH = 10,
NDAY = 25,
XTIME = 21600.

/
&NAM_PREP_WATFLUX  XTS_WATER_UNIF = 285.,
NYEAR = 2004,
NMONTH = 10,
NDAY = 25,
XTIME = 21600.

/
&NAM_PREP_TEB      XTI_ROAD= 285.,
XTI_BLD = 285.
XTS_ROAD= 285.
XTS_ROOF= 285.,
XTS_WALL= 285.,
XWS_ROAD= 0.,
XWS_ROOF= 0.,
NYEAR = 2004,
NMONTH = 10,
NDAY = 25,

```

```

/
XTIME = 21600.
/
&NAM_PREP_ISBA      XHUG_SURF = 0.2,
                    XHUG_ROOT = 0.2,
                    XHUG_DEEP = 0.2,
                    XTG_SURF  = 285.,
                    XTG_ROOT  = 288.,
                    XTG_DEEP  = 292.
                    NYEAR    = 2004,
                    NMONTH   = 10,
                    NDAY     = 25,
                    XTIME    = 21600.
/
&NAM_PREP_ISBA_SNOW  CSNOW = '3-L'
/

&NAM_IO_OFFLINE      YPROGRAM = 'ASCII' ,
                    YCOUPLING = 'E'
/
&NAM_DIAG_SURF_n     LSURF_BUDGET = T ,
                    N2M          = 1
/
&NAM_DIAG_SURF_ATM_n LFRAC      = T
/
&NAM_DIAG_ISBA_n     LPGD        = T ,
                    LSURF_EVAP_BUDGET = T ,
                    LSURF_MISC_BUDGET = T ,
                    LSURF_BUDGETC   = F
/
&NAM_DIAG_TEB_n      LSURF_MISC_BUDGET = T
/
&NAM_SGH_ISBA_n      CRUNOFF      = "WSAT"
/
&NAM_ISBA_n           CROUGH       = "Z04D" ,
                    CSCOND        = "NP89" ,
                    CALBEDO       = "DRY" ,
                    CC1DRY        = 'DEF' ,
                    CSOILFRZ      = 'DEF' ,
                    CDIFSFCOND    = 'DEF' ,
                    CSNOWRES      = 'DEF' ,
                    CCPSURF       = 'DRY'
/
&NAM_CH_ISBA_n        CCH_DRY_DEP = "WES89 "
/
&NAM_SEAFLUX_n        CSEA_ALB    = "TA96"
/
&NAM_CH_SEAFLUX_n     CCH_DRY_DEP = "WES89 "
/
&NAM_CH_WATFLUX_n     CCH_DRY_DEP = "WES89 "
/
&NAM_CH_TEB_n         CCH_DRY_DEP = "WES89 "
/

```

A.2.2 Uniform values prescribed : 1d example with patches

```

&NAM_DATA_ISBA      NTIME = 12 ,
                    XUNIF_VEGTYPE(1) = 0.,
                    XUNIF_VEGTYPE(2) = 0.,
                    XUNIF_VEGTYPE(3) = 0.,
                    XUNIF_VEGTYPE(4) = 0.,
                    XUNIF_VEGTYPE(5) = 0.,
                    XUNIF_VEGTYPE(6) = 0.,
                    XUNIF_VEGTYPE(7) = 1.,

```

XUNIF_VEGTYPE(8)	= 0.,				
XUNIF_VEGTYPE(9)	= 0.,				
XUNIF_VEGTYPE(10)	= 0.,				
XUNIF_VEGTYPE(11)	= 0.,				
XUNIF_VEGTYPE(12)	= 0.,				
XUNIF_VEG (1,1)	= 0.,	XUNIF_VEG (5,1)	= 0.,	XUNIF_VEG (9,1)	= 0.,
XUNIF_VEG (1,2)	= 0.,	XUNIF_VEG (5,2)	= 0.,	XUNIF_VEG (9,2)	= 0.,
XUNIF_VEG (1,3)	= 0.,	XUNIF_VEG (5,3)	= 0.,	XUNIF_VEG (9,3)	= 0.,
XUNIF_VEG (1,4)	= 0.,	XUNIF_VEG (5,4)	= 0.,	XUNIF_VEG (9,4)	= 0.,
XUNIF_VEG (1,5)	= 0.5,	XUNIF_VEG (5,5)	= 0.5,	XUNIF_VEG (9,5)	= 0.5,
XUNIF_VEG (1,6)	= 0.9,	XUNIF_VEG (5,6)	= 0.9,	XUNIF_VEG (9,6)	= 0.9,
XUNIF_VEG (1,7)	= 0.9,	XUNIF_VEG (5,7)	= 0.9,	XUNIF_VEG (9,7)	= 0.9,
XUNIF_VEG (1,8)	= 0.9,	XUNIF_VEG (5,8)	= 0.9,	XUNIF_VEG (9,8)	= 0.9,
XUNIF_VEG (1,9)	= 0.9,	XUNIF_VEG (5,9)	= 0.9,	XUNIF_VEG (9,9)	= 0.9,
XUNIF_VEG (1,10)	= 0.,	XUNIF_VEG (5,10)	= 0.,	XUNIF_VEG (9,10)	= 0.,
XUNIF_VEG (1,11)	= 0.,	XUNIF_VEG (5,11)	= 0.,	XUNIF_VEG (9,11)	= 0.,
XUNIF_VEG (1,12)	= 0.,	XUNIF_VEG (5,12)	= 0.,	XUNIF_VEG (9,12)	= 0.,
XUNIF_LAI (1,1)	= 0.,	XUNIF_LAI (5,1)	= 0.,	XUNIF_LAI (9,1)	= 0.,
XUNIF_LAI (1,2)	= 0.,	XUNIF_LAI (5,2)	= 0.,	XUNIF_LAI (9,2)	= 0.,
XUNIF_LAI (1,3)	= 0.,	XUNIF_LAI (5,3)	= 0.,	XUNIF_LAI (9,3)	= 0.,
XUNIF_LAI (1,4)	= 0.,	XUNIF_LAI (5,4)	= 0.,	XUNIF_LAI (9,4)	= 0.,
XUNIF_LAI (1,5)	= 1.,	XUNIF_LAI (5,5)	= 1.,	XUNIF_LAI (9,5)	= 1.,
XUNIF_LAI (1,6)	= 3.,	XUNIF_LAI (5,6)	= 3.,	XUNIF_LAI (9,6)	= 3.,
XUNIF_LAI (1,7)	= 3.,	XUNIF_LAI (5,7)	= 3.,	XUNIF_LAI (9,7)	= 3.,
XUNIF_LAI (1,8)	= 3.,	XUNIF_LAI (5,8)	= 3.,	XUNIF_LAI (9,8)	= 3.,
XUNIF_LAI (1,9)	= 3.,	XUNIF_LAI (5,9)	= 3.,	XUNIF_LAI (9,9)	= 3.,
XUNIF_LAI (1,10)	= 0.,	XUNIF_LAI (5,10)	= 0.,	XUNIF_LAI (9,10)	= 0.,
XUNIF_LAI (1,11)	= 0.,	XUNIF_LAI (5,11)	= 0.,	XUNIF_LAI (9,11)	= 0.,
XUNIF_LAI (1,12)	= 0.,	XUNIF_LAI (5,12)	= 0.,	XUNIF_LAI (9,12)	= 0.,
XUNIF_ZO (1,1)	= 0.01,	XUNIF_ZO (5,1)	= 0.01,	XUNIF_ZO (9,1)	= 0.01,
XUNIF_ZO (1,2)	= 0.01,	XUNIF_ZO (5,2)	= 0.01,	XUNIF_ZO (9,2)	= 0.01,
XUNIF_ZO (1,3)	= 0.01,	XUNIF_ZO (5,3)	= 0.01,	XUNIF_ZO (9,3)	= 0.01,
XUNIF_ZO (1,4)	= 0.01,	XUNIF_ZO (5,4)	= 0.01,	XUNIF_ZO (9,4)	= 0.01,
XUNIF_ZO (1,5)	= 0.05,	XUNIF_ZO (5,5)	= 0.05,	XUNIF_ZO (9,5)	= 0.05,
XUNIF_ZO (1,6)	= 0.15,	XUNIF_ZO (5,6)	= 0.15,	XUNIF_ZO (9,6)	= 0.15,
XUNIF_ZO (1,7)	= 0.15,	XUNIF_ZO (5,7)	= 0.15,	XUNIF_ZO (9,7)	= 0.15,
XUNIF_ZO (1,8)	= 0.15,	XUNIF_ZO (5,8)	= 0.15,	XUNIF_ZO (9,8)	= 0.15,
XUNIF_ZO (1,9)	= 0.15,	XUNIF_ZO (5,9)	= 0.15,	XUNIF_ZO (9,9)	= 0.15,
XUNIF_ZO (1,10)	= 0.01,	XUNIF_ZO (5,10)	= 0.01,	XUNIF_ZO (9,10)	= 0.01,
XUNIF_ZO (1,11)	= 0.01,	XUNIF_ZO (5,11)	= 0.01,	XUNIF_ZO (9,11)	= 0.01,
XUNIF_ZO (1,12)	= 0.01,	XUNIF_ZO (5,12)	= 0.01,	XUNIF_ZO (9,12)	= 0.01,
XUNIF_EMIS(1,1)	= 0.98,	XUNIF_EMIS(5,1)	= 0.98,	XUNIF_EMIS(9,1)	= 0.98,
XUNIF_EMIS(1,2)	= 0.98,	XUNIF_EMIS(5,2)	= 0.98,	XUNIF_EMIS(9,2)	= 0.98,
XUNIF_EMIS(1,3)	= 0.98,	XUNIF_EMIS(5,3)	= 0.98,	XUNIF_EMIS(9,3)	= 0.98,
XUNIF_EMIS(1,4)	= 0.98,	XUNIF_EMIS(5,4)	= 0.98,	XUNIF_EMIS(9,4)	= 0.98,
XUNIF_EMIS(1,5)	= 0.98,	XUNIF_EMIS(5,5)	= 0.98,	XUNIF_EMIS(9,5)	= 0.98,
XUNIF_EMIS(1,6)	= 0.98,	XUNIF_EMIS(5,6)	= 0.98,	XUNIF_EMIS(9,6)	= 0.98,
XUNIF_EMIS(1,7)	= 0.98,	XUNIF_EMIS(5,7)	= 0.98,	XUNIF_EMIS(9,7)	= 0.98,
XUNIF_EMIS(1,8)	= 0.98,	XUNIF_EMIS(5,8)	= 0.98,	XUNIF_EMIS(9,8)	= 0.98,
XUNIF_EMIS(1,9)	= 0.98,	XUNIF_EMIS(5,9)	= 0.98,	XUNIF_EMIS(9,9)	= 0.98,
XUNIF_EMIS(1,10)	= 0.98,	XUNIF_EMIS(5,10)	= 0.98,	XUNIF_EMIS(9,10)	= 0.98,
XUNIF_EMIS(1,11)	= 0.98,	XUNIF_EMIS(5,11)	= 0.98,	XUNIF_EMIS(9,11)	= 0.98,
XUNIF_EMIS(1,12)	= 0.98,	XUNIF_EMIS(5,12)	= 0.98,	XUNIF_EMIS(9,12)	= 0.98,
XUNIF_VEG (2,1)	= 0.,	XUNIF_VEG (6,1)	= 0.,	XUNIF_VEG (10,1)	= 0.,
XUNIF_VEG (2,2)	= 0.,	XUNIF_VEG (6,2)	= 0.,	XUNIF_VEG (10,2)	= 0.,
XUNIF_VEG (2,3)	= 0.,	XUNIF_VEG (6,3)	= 0.,	XUNIF_VEG (10,3)	= 0.,
XUNIF_VEG (2,4)	= 0.,	XUNIF_VEG (6,4)	= 0.,	XUNIF_VEG (10,4)	= 0.,
XUNIF_VEG (2,5)	= 0.5,	XUNIF_VEG (6,5)	= 0.5,	XUNIF_VEG (10,5)	= 0.5,
XUNIF_VEG (2,6)	= 0.9,	XUNIF_VEG (6,6)	= 0.9,	XUNIF_VEG (10,6)	= 0.9,
XUNIF_VEG (2,7)	= 0.9,	XUNIF_VEG (6,7)	= 0.9,	XUNIF_VEG (10,7)	= 0.9,
XUNIF_VEG (2,8)	= 0.9,	XUNIF_VEG (6,8)	= 0.9,	XUNIF_VEG (10,8)	= 0.9,
XUNIF_VEG (2,9)	= 0.9,	XUNIF_VEG (6,9)	= 0.9,	XUNIF_VEG (10,9)	= 0.9,
XUNIF_VEG (2,10)	= 0.,	XUNIF_VEG (6,10)	= 0.,	XUNIF_VEG (10,10)	= 0.,
XUNIF_VEG (2,11)	= 0.,	XUNIF_VEG (6,11)	= 0.,	XUNIF_VEG (10,11)	= 0.,
XUNIF_VEG (2,12)	= 0.,	XUNIF_VEG (6,12)	= 0.,	XUNIF_VEG (10,12)	= 0.,
XUNIF_LAI (2,1)	= 0.,	XUNIF_LAI (6,1)	= 0.,	XUNIF_LAI (10,1)	= 0.,
XUNIF_LAI (2,2)	= 0.,	XUNIF_LAI (6,2)	= 0.,	XUNIF_LAI (10,2)	= 0.,
XUNIF_LAI (2,3)	= 0.,	XUNIF_LAI (6,3)	= 0.,	XUNIF_LAI (10,3)	= 0.,

XUNIF_LAI (2,4)	= 0.,	XUNIF_LAI (6,4)	= 0.,	XUNIF_LAI (10,4)	= 0.,
XUNIF_LAI (2,5)	= 1.,	XUNIF_LAI (6,5)	= 1.,	XUNIF_LAI (10,5)	= 1.,
XUNIF_LAI (2,6)	= 3.,	XUNIF_LAI (6,6)	= 3.,	XUNIF_LAI (10,6)	= 3.,
XUNIF_LAI (2,7)	= 3.,	XUNIF_LAI (6,7)	= 3.,	XUNIF_LAI (10,7)	= 3.,
XUNIF_LAI (2,8)	= 3.,	XUNIF_LAI (6,8)	= 3.,	XUNIF_LAI (10,8)	= 3.,
XUNIF_LAI (2,9)	= 3.,	XUNIF_LAI (6,9)	= 3.,	XUNIF_LAI (10,9)	= 3.,
XUNIF_LAI (2,10)	= 0.,	XUNIF_LAI (6,10)	= 0.,	XUNIF_LAI (10,10)	= 0.,
XUNIF_LAI (2,11)	= 0.,	XUNIF_LAI (6,11)	= 0.,	XUNIF_LAI (10,11)	= 0.,
XUNIF_LAI (2,12)	= 0.,	XUNIF_LAI (6,12)	= 0.,	XUNIF_LAI (10,12)	= 0.,
XUNIF_ZO (2,1)	= 0.01,	XUNIF_ZO (6,1)	= 0.01,	XUNIF_ZO (10,1)	= 0.01,
XUNIF_ZO (2,2)	= 0.01,	XUNIF_ZO (6,2)	= 0.01,	XUNIF_ZO (10,2)	= 0.01,
XUNIF_ZO (2,3)	= 0.01,	XUNIF_ZO (6,3)	= 0.01,	XUNIF_ZO (10,3)	= 0.01,
XUNIF_ZO (2,4)	= 0.01,	XUNIF_ZO (6,4)	= 0.01,	XUNIF_ZO (10,4)	= 0.01,
XUNIF_ZO (2,5)	= 0.05,	XUNIF_ZO (6,5)	= 0.05,	XUNIF_ZO (10,5)	= 0.05,
XUNIF_ZO (2,6)	= 0.15,	XUNIF_ZO (6,6)	= 0.15,	XUNIF_ZO (10,6)	= 0.15,
XUNIF_ZO (2,7)	= 0.15,	XUNIF_ZO (6,7)	= 0.15,	XUNIF_ZO (10,7)	= 0.15,
XUNIF_ZO (2,8)	= 0.15,	XUNIF_ZO (6,8)	= 0.15,	XUNIF_ZO (10,8)	= 0.15,
XUNIF_ZO (2,9)	= 0.15,	XUNIF_ZO (6,9)	= 0.15,	XUNIF_ZO (10,9)	= 0.15,
XUNIF_ZO (2,10)	= 0.01,	XUNIF_ZO (6,10)	= 0.01,	XUNIF_ZO (10,10)	= 0.01,
XUNIF_ZO (2,11)	= 0.01,	XUNIF_ZO (6,11)	= 0.01,	XUNIF_ZO (10,11)	= 0.01,
XUNIF_ZO (2,12)	= 0.01,	XUNIF_ZO (6,12)	= 0.01,	XUNIF_ZO (10,12)	= 0.01,
XUNIF_EMIS(2,1)	= 0.98,	XUNIF_EMIS(6,1)	= 0.98,	XUNIF_EMIS(10,1)	= 0.98,
XUNIF_EMIS(2,2)	= 0.98,	XUNIF_EMIS(6,2)	= 0.98,	XUNIF_EMIS(10,2)	= 0.98,
XUNIF_EMIS(2,3)	= 0.98,	XUNIF_EMIS(6,3)	= 0.98,	XUNIF_EMIS(10,3)	= 0.98,
XUNIF_EMIS(2,4)	= 0.98,	XUNIF_EMIS(6,4)	= 0.98,	XUNIF_EMIS(10,4)	= 0.98,
XUNIF_EMIS(2,5)	= 0.98,	XUNIF_EMIS(6,5)	= 0.98,	XUNIF_EMIS(10,5)	= 0.98,
XUNIF_EMIS(2,6)	= 0.98,	XUNIF_EMIS(6,6)	= 0.98,	XUNIF_EMIS(10,6)	= 0.98,
XUNIF_EMIS(2,7)	= 0.98,	XUNIF_EMIS(6,7)	= 0.98,	XUNIF_EMIS(10,7)	= 0.98,
XUNIF_EMIS(2,8)	= 0.98,	XUNIF_EMIS(6,8)	= 0.98,	XUNIF_EMIS(10,8)	= 0.98,
XUNIF_EMIS(2,9)	= 0.98,	XUNIF_EMIS(6,9)	= 0.98,	XUNIF_EMIS(10,9)	= 0.98,
XUNIF_EMIS(2,10)	= 0.98,	XUNIF_EMIS(6,10)	= 0.98,	XUNIF_EMIS(10,10)	= 0.98,
XUNIF_EMIS(2,11)	= 0.98,	XUNIF_EMIS(6,11)	= 0.98,	XUNIF_EMIS(10,11)	= 0.98,
XUNIF_EMIS(2,12)	= 0.98,	XUNIF_EMIS(6,12)	= 0.98,	XUNIF_EMIS(10,12)	= 0.98,
XUNIF_VEG (3,1)	= 0.,	XUNIF_VEG (7,1)	= 0.,	XUNIF_VEG (11,1)	= 0.,
XUNIF_VEG (3,2)	= 0.,	XUNIF_VEG (7,2)	= 0.,	XUNIF_VEG (11,2)	= 0.,
XUNIF_VEG (3,3)	= 0.,	XUNIF_VEG (7,3)	= 0.,	XUNIF_VEG (11,3)	= 0.,
XUNIF_VEG (3,4)	= 0.,	XUNIF_VEG (7,4)	= 0.,	XUNIF_VEG (11,4)	= 0.,
XUNIF_VEG (3,5)	= 0.5,	XUNIF_VEG (7,5)	= 0.5,	XUNIF_VEG (11,5)	= 0.5,
XUNIF_VEG (3,6)	= 0.9,	XUNIF_VEG (7,6)	= 0.9,	XUNIF_VEG (11,6)	= 0.9,
XUNIF_VEG (3,7)	= 0.9,	XUNIF_VEG (7,7)	= 0.9,	XUNIF_VEG (11,7)	= 0.9,
XUNIF_VEG (3,8)	= 0.9,	XUNIF_VEG (7,8)	= 0.9,	XUNIF_VEG (11,8)	= 0.9,
XUNIF_VEG (3,9)	= 0.9,	XUNIF_VEG (7,9)	= 0.9,	XUNIF_VEG (11,9)	= 0.9,
XUNIF_VEG (3,10)	= 0.,	XUNIF_VEG (7,10)	= 0.,	XUNIF_VEG (11,10)	= 0.,
XUNIF_VEG (3,11)	= 0.,	XUNIF_VEG (7,11)	= 0.,	XUNIF_VEG (11,11)	= 0.,
XUNIF_VEG (3,12)	= 0.,	XUNIF_VEG (7,12)	= 0.,	XUNIF_VEG (11,12)	= 0.,
XUNIF_LAI (3,1)	= 0.,	XUNIF_LAI (7,1)	= 0.,	XUNIF_LAI (11,1)	= 0.,
XUNIF_LAI (3,2)	= 0.,	XUNIF_LAI (7,2)	= 0.,	XUNIF_LAI (11,2)	= 0.,
XUNIF_LAI (3,3)	= 0.,	XUNIF_LAI (7,3)	= 0.,	XUNIF_LAI (11,3)	= 0.,
XUNIF_LAI (3,4)	= 0.,	XUNIF_LAI (7,4)	= 0.,	XUNIF_LAI (11,4)	= 0.,
XUNIF_LAI (3,5)	= 1.,	XUNIF_LAI (7,5)	= 1.,	XUNIF_LAI (11,5)	= 1.,
XUNIF_LAI (3,6)	= 3.,	XUNIF_LAI (7,6)	= 3.,	XUNIF_LAI (11,6)	= 3.,
XUNIF_LAI (3,7)	= 3.,	XUNIF_LAI (7,7)	= 3.,	XUNIF_LAI (11,7)	= 3.,
XUNIF_LAI (3,8)	= 3.,	XUNIF_LAI (7,8)	= 3.,	XUNIF_LAI (11,8)	= 3.,
XUNIF_LAI (3,9)	= 3.,	XUNIF_LAI (7,9)	= 3.,	XUNIF_LAI (11,9)	= 3.,
XUNIF_LAI (3,10)	= 0.,	XUNIF_LAI (7,10)	= 0.,	XUNIF_LAI (11,10)	= 0.,
XUNIF_LAI (3,11)	= 0.,	XUNIF_LAI (7,11)	= 0.,	XUNIF_LAI (11,11)	= 0.,
XUNIF_LAI (3,12)	= 0.,	XUNIF_LAI (7,12)	= 0.,	XUNIF_LAI (11,12)	= 0.,
XUNIF_ZO (3,1)	= 0.01,	XUNIF_ZO (7,1)	= 0.01,	XUNIF_ZO (11,1)	= 0.01,
XUNIF_ZO (3,2)	= 0.01,	XUNIF_ZO (7,2)	= 0.01,	XUNIF_ZO (11,2)	= 0.01,
XUNIF_ZO (3,3)	= 0.01,	XUNIF_ZO (7,3)	= 0.01,	XUNIF_ZO (11,3)	= 0.01,
XUNIF_ZO (3,4)	= 0.01,	XUNIF_ZO (7,4)	= 0.01,	XUNIF_ZO (11,4)	= 0.01,
XUNIF_ZO (3,5)	= 0.05,	XUNIF_ZO (7,5)	= 0.05,	XUNIF_ZO (11,5)	= 0.05,
XUNIF_ZO (3,6)	= 0.15,	XUNIF_ZO (7,6)	= 0.15,	XUNIF_ZO (11,6)	= 0.15,
XUNIF_ZO (3,7)	= 0.15,	XUNIF_ZO (7,7)	= 0.15,	XUNIF_ZO (11,7)	= 0.15,
XUNIF_ZO (3,8)	= 0.15,	XUNIF_ZO (7,8)	= 0.15,	XUNIF_ZO (11,8)	= 0.15,
XUNIF_ZO (3,9)	= 0.15,	XUNIF_ZO (7,9)	= 0.15,	XUNIF_ZO (11,9)	= 0.15,
XUNIF_ZO (3,10)	= 0.01,	XUNIF_ZO (7,10)	= 0.01,	XUNIF_ZO (11,10)	= 0.01,
XUNIF_ZO (3,11)	= 0.01,	XUNIF_ZO (7,11)	= 0.01,	XUNIF_ZO (11,11)	= 0.01,

XUNIF_ZO (3,12)	= 0.01,	XUNIF_ZO (7,12)	= 0.01,	XUNIF_ZO (11,12)	= 0.01,
XUNIF_EMIS(3,1)	= 0.98,	XUNIF_EMIS(7,1)	= 0.98,	XUNIF_EMIS(11,1)	= 0.98,
XUNIF_EMIS(3,2)	= 0.98,	XUNIF_EMIS(7,2)	= 0.98,	XUNIF_EMIS(11,2)	= 0.98,
XUNIF_EMIS(3,3)	= 0.98,	XUNIF_EMIS(7,3)	= 0.98,	XUNIF_EMIS(11,3)	= 0.98,
XUNIF_EMIS(3,4)	= 0.98,	XUNIF_EMIS(7,4)	= 0.98,	XUNIF_EMIS(11,4)	= 0.98,
XUNIF_EMIS(3,5)	= 0.98,	XUNIF_EMIS(7,5)	= 0.98,	XUNIF_EMIS(11,5)	= 0.98,
XUNIF_EMIS(3,6)	= 0.98,	XUNIF_EMIS(7,6)	= 0.98,	XUNIF_EMIS(11,6)	= 0.98,
XUNIF_EMIS(3,7)	= 0.98,	XUNIF_EMIS(7,7)	= 0.98,	XUNIF_EMIS(11,7)	= 0.98,
XUNIF_EMIS(3,8)	= 0.98,	XUNIF_EMIS(7,8)	= 0.98,	XUNIF_EMIS(11,8)	= 0.98,
XUNIF_EMIS(3,9)	= 0.98,	XUNIF_EMIS(7,9)	= 0.98,	XUNIF_EMIS(11,9)	= 0.98,
XUNIF_EMIS(3,10)	= 0.98,	XUNIF_EMIS(7,10)	= 0.98,	XUNIF_EMIS(11,10)	= 0.98,
XUNIF_EMIS(3,11)	= 0.98,	XUNIF_EMIS(7,11)	= 0.98,	XUNIF_EMIS(11,11)	= 0.98,
XUNIF_EMIS(3,12)	= 0.98,	XUNIF_EMIS(7,12)	= 0.98,	XUNIF_EMIS(11,12)	= 0.98,
XUNIF_VEG (4,1)	= 0.,	XUNIF_VEG (8,1)	= 0.,	XUNIF_VEG (12,1)	= 0.,
XUNIF_VEG (4,2)	= 0.,	XUNIF_VEG (8,2)	= 0.,	XUNIF_VEG (12,2)	= 0.,
XUNIF_VEG (4,3)	= 0.,	XUNIF_VEG (8,3)	= 0.,	XUNIF_VEG (12,3)	= 0.,
XUNIF_VEG (4,4)	= 0.,	XUNIF_VEG (8,4)	= 0.,	XUNIF_VEG (12,4)	= 0.,
XUNIF_VEG (4,5)	= 0.5,	XUNIF_VEG (8,5)	= 0.5,	XUNIF_VEG (12,5)	= 0.5,
XUNIF_VEG (4,6)	= 0.9,	XUNIF_VEG (8,6)	= 0.9,	XUNIF_VEG (12,6)	= 0.9,
XUNIF_VEG (4,7)	= 0.9,	XUNIF_VEG (8,7)	= 0.9,	XUNIF_VEG (12,7)	= 0.9,
XUNIF_VEG (4,8)	= 0.9,	XUNIF_VEG (8,8)	= 0.9,	XUNIF_VEG (12,8)	= 0.9,
XUNIF_VEG (4,9)	= 0.9,	XUNIF_VEG (8,9)	= 0.9,	XUNIF_VEG (12,9)	= 0.9,
XUNIF_VEG (4,10)	= 0.,	XUNIF_VEG (8,10)	= 0.,	XUNIF_VEG (12,10)	= 0.,
XUNIF_VEG (4,11)	= 0.,	XUNIF_VEG (8,11)	= 0.,	XUNIF_VEG (12,11)	= 0.,
XUNIF_VEG (4,12)	= 0.,	XUNIF_VEG (8,12)	= 0.,	XUNIF_VEG (12,12)	= 0.,
XUNIF_LAI (4,1)	= 0.,	XUNIF_LAI (8,1)	= 0.,	XUNIF_LAI (12,1)	= 0.,
XUNIF_LAI (4,2)	= 0.,	XUNIF_LAI (8,2)	= 0.,	XUNIF_LAI (12,2)	= 0.,
XUNIF_LAI (4,3)	= 0.,	XUNIF_LAI (8,3)	= 0.,	XUNIF_LAI (12,3)	= 0.,
XUNIF_LAI (4,4)	= 0.,	XUNIF_LAI (8,4)	= 0.,	XUNIF_LAI (12,4)	= 0.,
XUNIF_LAI (4,5)	= 1.,	XUNIF_LAI (8,5)	= 1.,	XUNIF_LAI (12,5)	= 1.,
XUNIF_LAI (4,6)	= 3.,	XUNIF_LAI (8,6)	= 3.,	XUNIF_LAI (12,6)	= 3.,
XUNIF_LAI (4,7)	= 3.,	XUNIF_LAI (8,7)	= 3.,	XUNIF_LAI (12,7)	= 3.,
XUNIF_LAI (4,8)	= 3.,	XUNIF_LAI (8,8)	= 3.,	XUNIF_LAI (12,8)	= 3.,
XUNIF_LAI (4,9)	= 3.,	XUNIF_LAI (8,9)	= 3.,	XUNIF_LAI (12,9)	= 3.,
XUNIF_LAI (4,10)	= 0.,	XUNIF_LAI (8,10)	= 0.,	XUNIF_LAI (12,10)	= 0.,
XUNIF_LAI (4,11)	= 0.,	XUNIF_LAI (8,11)	= 0.,	XUNIF_LAI (12,11)	= 0.,
XUNIF_LAI (4,12)	= 0.,	XUNIF_LAI (8,12)	= 0.,	XUNIF_LAI (12,12)	= 0.,
XUNIF_ZO (4,1)	= 0.01,	XUNIF_ZO (8,1)	= 0.01,	XUNIF_ZO (12,1)	= 0.01,
XUNIF_ZO (4,2)	= 0.01,	XUNIF_ZO (8,2)	= 0.01,	XUNIF_ZO (12,2)	= 0.01,
XUNIF_ZO (4,3)	= 0.01,	XUNIF_ZO (8,3)	= 0.01,	XUNIF_ZO (12,3)	= 0.01,
XUNIF_ZO (4,4)	= 0.01,	XUNIF_ZO (8,4)	= 0.01,	XUNIF_ZO (12,4)	= 0.01,
XUNIF_ZO (4,5)	= 0.05,	XUNIF_ZO (8,5)	= 0.05,	XUNIF_ZO (12,5)	= 0.05,
XUNIF_ZO (4,6)	= 0.15,	XUNIF_ZO (8,6)	= 0.15,	XUNIF_ZO (12,6)	= 0.15,
XUNIF_ZO (4,7)	= 0.15,	XUNIF_ZO (8,7)	= 0.15,	XUNIF_ZO (12,7)	= 0.15,
XUNIF_ZO (4,8)	= 0.15,	XUNIF_ZO (8,8)	= 0.15,	XUNIF_ZO (12,8)	= 0.15,
XUNIF_ZO (4,9)	= 0.15,	XUNIF_ZO (8,9)	= 0.15,	XUNIF_ZO (12,9)	= 0.15,
XUNIF_ZO (4,10)	= 0.01,	XUNIF_ZO (8,10)	= 0.01,	XUNIF_ZO (12,10)	= 0.01,
XUNIF_ZO (4,11)	= 0.01,	XUNIF_ZO (8,11)	= 0.01,	XUNIF_ZO (12,11)	= 0.01,
XUNIF_ZO (4,12)	= 0.01,	XUNIF_ZO (8,12)	= 0.01,	XUNIF_ZO (12,12)	= 0.01,
XUNIF_EMIS(4,1)	= 0.98,	XUNIF_EMIS(8,1)	= 0.98,	XUNIF_EMIS(12,1)	= 0.98,
XUNIF_EMIS(4,2)	= 0.98,	XUNIF_EMIS(8,2)	= 0.98,	XUNIF_EMIS(12,2)	= 0.98,
XUNIF_EMIS(4,3)	= 0.98,	XUNIF_EMIS(8,3)	= 0.98,	XUNIF_EMIS(12,3)	= 0.98,
XUNIF_EMIS(4,4)	= 0.98,	XUNIF_EMIS(8,4)	= 0.98,	XUNIF_EMIS(12,4)	= 0.98,
XUNIF_EMIS(4,5)	= 0.98,	XUNIF_EMIS(8,5)	= 0.98,	XUNIF_EMIS(12,5)	= 0.98,
XUNIF_EMIS(4,6)	= 0.98,	XUNIF_EMIS(8,6)	= 0.98,	XUNIF_EMIS(12,6)	= 0.98,
XUNIF_EMIS(4,7)	= 0.98,	XUNIF_EMIS(8,7)	= 0.98,	XUNIF_EMIS(12,7)	= 0.98,
XUNIF_EMIS(4,8)	= 0.98,	XUNIF_EMIS(8,8)	= 0.98,	XUNIF_EMIS(12,8)	= 0.98,
XUNIF_EMIS(4,9)	= 0.98,	XUNIF_EMIS(8,9)	= 0.98,	XUNIF_EMIS(12,9)	= 0.98,
XUNIF_EMIS(4,10)	= 0.98,	XUNIF_EMIS(8,10)	= 0.98,	XUNIF_EMIS(12,10)	= 0.98,
XUNIF_EMIS(4,11)	= 0.98,	XUNIF_EMIS(8,11)	= 0.98,	XUNIF_EMIS(12,11)	= 0.98,
XUNIF_EMIS(4,12)	= 0.98,	XUNIF_EMIS(8,12)	= 0.98,	XUNIF_EMIS(12,12)	= 0.98,
XUNIF_DG(1,1)	= 0.01,	XUNIF_DG(5,1)	= 0.01,	XUNIF_DG(9,1)	= 0.01,
XUNIF_DG(1,2)	= 1.60,	XUNIF_DG(5,2)	= 1.60,	XUNIF_DG(9,2)	= 1.60,
XUNIF_DG(1,3)	= 1.60,	XUNIF_DG(5,3)	= 1.60,	XUNIF_DG(9,3)	= 1.60,
XUNIF_DG(2,1)	= 0.01,	XUNIF_DG(6,1)	= 0.01,	XUNIF_DG(10,1)	= 0.01,
XUNIF_DG(2,2)	= 1.60,	XUNIF_DG(6,2)	= 1.60,	XUNIF_DG(10,2)	= 1.60,
XUNIF_DG(2,3)	= 1.60,	XUNIF_DG(6,3)	= 1.60,	XUNIF_DG(10,3)	= 1.60,
XUNIF_DG(3,1)	= 0.01,	XUNIF_DG(7,1)	= 0.01,	XUNIF_DG(11,1)	= 0.01,

XUNIF_DG(3,2)	= 1.60,	XUNIF_DG(7,2)	= 1.60,	XUNIF_DG(11,2)	= 1.60,
XUNIF_DG(3,3)	= 1.60,	XUNIF_DG(7,3)	= 1.60,	XUNIF_DG(11,3)	= 1.60,
XUNIF_DG(4,1)	= 0.01,	XUNIF_DG(8,1)	= 0.01,	XUNIF_DG(12,1)	= 0.01,
XUNIF_DG(4,2)	= 1.60,	XUNIF_DG(8,2)	= 1.60,	XUNIF_DG(12,2)	= 1.60,
XUNIF_DG(4,3)	= 1.60,	XUNIF_DG(8,3)	= 1.60,	XUNIF_DG(12,3)	= 1.60,
XUNIF_ROOTFRAC(1,1)	= -999.,	XUNIF_ROOTFRAC(5,1)	= -999.,	XUNIF_ROOTFRAC(9,1)	= -999.,
XUNIF_ROOTFRAC(1,2)	= -999.,	XUNIF_ROOTFRAC(5,2)	= -999.,	XUNIF_ROOTFRAC(9,2)	= -999.,
XUNIF_ROOTFRAC(1,3)	= -999.,	XUNIF_ROOTFRAC(5,3)	= -999.,	XUNIF_ROOTFRAC(9,3)	= -999.,
XUNIF_ROOTFRAC(2,1)	= -999.,	XUNIF_ROOTFRAC(6,1)	= -999.,	XUNIF_ROOTFRAC(10,1)	= -999.,
XUNIF_ROOTFRAC(2,2)	= -999.,	XUNIF_ROOTFRAC(6,2)	= -999.,	XUNIF_ROOTFRAC(10,2)	= -999.,
XUNIF_ROOTFRAC(2,3)	= -999.,	XUNIF_ROOTFRAC(6,3)	= -999.,	XUNIF_ROOTFRAC(10,3)	= -999.,
XUNIF_ROOTFRAC(3,1)	= -999.,	XUNIF_ROOTFRAC(7,1)	= -999.,	XUNIF_ROOTFRAC(11,1)	= -999.,
XUNIF_ROOTFRAC(3,2)	= -999.,	XUNIF_ROOTFRAC(7,2)	= -999.,	XUNIF_ROOTFRAC(11,2)	= -999.,
XUNIF_ROOTFRAC(3,3)	= -999.,	XUNIF_ROOTFRAC(7,3)	= -999.,	XUNIF_ROOTFRAC(11,3)	= -999.,
XUNIF_ROOTFRAC(4,1)	= -999.,	XUNIF_ROOTFRAC(8,1)	= -999.,	XUNIF_ROOTFRAC(12,1)	= -999.,
XUNIF_ROOTFRAC(4,2)	= -999.,	XUNIF_ROOTFRAC(8,2)	= -999.,	XUNIF_ROOTFRAC(12,2)	= -999.,
XUNIF_ROOTFRAC(4,3)	= -999.,	XUNIF_ROOTFRAC(8,3)	= -999.,	XUNIF_ROOTFRAC(12,3)	= -999.,
XUNIF_RSMIN	(1)= 40.,	XUNIF_RSMIN	(2)= 40.,	XUNIF_RSMIN	(3)= 40.,
XUNIF_GAMMA	(1)= 0.,	XUNIF_GAMMA	(2)= 0.,	XUNIF_GAMMA	(3)= 0.,
XUNIF_WRMAX_CF	(1)= 0.2,	XUNIF_WRMAX_CF	(2)= 0.2,	XUNIF_WRMAX_CF	(3)= 0.2,
XUNIF_RGL	(1)= 100.,	XUNIF_RGL	(2)= 100.,	XUNIF_RGL	(3)= 100.,
XUNIF_CV	(1)= 0.00002,	XUNIF_CV	(2)= 0.00002,	XUNIF_CV	(3)= 0.00002,
XUNIF_ZO_O_ZOH	(1)= 10.,	XUNIF_ZO_O_ZOH	(2)= 10.,	XUNIF_ZO_O_ZOH	(3)= 10.,
XUNIF_ALBNIR_VEG	(1)= 0.3,	XUNIF_ALBNIR_VEG	(2)= 0.3,	XUNIF_ALBNIR_VEG	(3)= 0.3,
XUNIF_ALBVIS_VEG	(1)= 0.1,	XUNIF_ALBVIS_VEG	(2)= 0.1,	XUNIF_ALBVIS_VEG	(3)= 0.1,
XUNIF_ALBUV_VEG	(1)= 0.0425,	XUNIF_ALBUV_VEG	(2)= 0.0425,	XUNIF_ALBUV_VEG	(3)= 0.0425,
XUNIF_ALBNIR_SOIL	(1)= 0.3,	XUNIF_ALBNIR_SOIL	(2)= 0.3,	XUNIF_ALBNIR_SOIL	(3)= 0.3,
XUNIF_ALBVIS_SOIL	(1)= 0.1,	XUNIF_ALBVIS_SOIL	(2)= 0.1,	XUNIF_ALBVIS_SOIL	(3)= 0.1,
XUNIF_ALBUV_SOIL	(1)= 0.06,	XUNIF_ALBUV_SOIL	(2)= 0.06,	XUNIF_ALBUV_SOIL	(3)= 0.06,
XUNIF_GMES	(1)= 0.003,	XUNIF_GMES	(2)= 0.003,	XUNIF_GMES	(3)= 0.003,
XUNIF_RE25	(1)= 0.0000003,	XUNIF_RE25	(2)= 0.0000003,	XUNIF_RE25	(3)= 0.0000003,
XUNIF_BSLAI	(1)= 0.06,	XUNIF_BSLAI	(2)= 0.06,	XUNIF_BSLAI	(3)= 0.06,
XUNIF_LAIMIN	(1)= 0.3,	XUNIF_LAIMIN	(2)= 0.3,	XUNIF_LAIMIN	(3)= 0.3,
XUNIF_SEFOLD	(1)= 5184000.,	XUNIF_SEFOLD	(2)= 5184000.,	XUNIF_SEFOLD	(3)= 5184000.,
XUNIF_GC	(1)= 0.00025,	XUNIF_GC	(2)= 0.00025,	XUNIF_GC	(3)= 0.00025,
XUNIF_DMAX	(1)= 0.1,	XUNIF_DMAX	(2)= 0.1,	XUNIF_DMAX	(3)= 0.1,
XUNIF_F2I	(1)= 0.3,	XUNIF_F2I	(2)= 0.3,	XUNIF_F2I	(3)= 0.3,
XUNIF_H_TREE	(1)= 20.,	XUNIF_H_TREE	(2)= 20.,	XUNIF_H_TREE	(3)= 20.,
XUNIF_CE_NITRO	(1)= 3.79,	XUNIF_CE_NITRO	(2)= 3.79,	XUNIF_CE_NITRO	(3)= 3.79,
XUNIF_CF_NITRO	(1)= 9.84,	XUNIF_CF_NITRO	(2)= 9.84,	XUNIF_CF_NITRO	(3)= 9.84,
XUNIF_CNA_NITRO	(1)= 1.3,	XUNIF_CNA_NITRO	(2)= 1.3,	XUNIF_CNA_NITRO	(3)= 1.3,
XUNIF_RSMIN	(4)= 40.,	XUNIF_RSMIN	(5)= 40.,	XUNIF_RSMIN	(6)= 40.,
XUNIF_GAMMA	(4)= 0.,	XUNIF_GAMMA	(5)= 0.,	XUNIF_GAMMA	(6)= 0.,
XUNIF_WRMAX_CF	(4)= 0.2,	XUNIF_WRMAX_CF	(5)= 0.2,	XUNIF_WRMAX_CF	(6)= 0.2,
XUNIF_RGL	(4)= 100.,	XUNIF_RGL	(5)= 100.,	XUNIF_RGL	(6)= 100.,
XUNIF_CV	(4)= 0.00002,	XUNIF_CV	(5)= 0.00002,	XUNIF_CV	(6)= 0.00002,
XUNIF_ZO_O_ZOH	(4)= 10.,	XUNIF_ZO_O_ZOH	(5)= 10.,	XUNIF_ZO_O_ZOH	(6)= 10.,
XUNIF_ALBNIR_VEG	(4)= 0.3,	XUNIF_ALBNIR_VEG	(5)= 0.3,	XUNIF_ALBNIR_VEG	(6)= 0.3,
XUNIF_ALBVIS_VEG	(4)= 0.1,	XUNIF_ALBVIS_VEG	(5)= 0.1,	XUNIF_ALBVIS_VEG	(6)= 0.1,
XUNIF_ALBUV_VEG	(4)= 0.0425,	XUNIF_ALBUV_VEG	(5)= 0.0425,	XUNIF_ALBUV_VEG	(6)= 0.0425,
XUNIF_ALBNIR_SOIL	(4)= 0.3,	XUNIF_ALBNIR_SOIL	(5)= 0.3,	XUNIF_ALBNIR_SOIL	(6)= 0.3,
XUNIF_ALBVIS_SOIL	(4)= 0.1,	XUNIF_ALBVIS_SOIL	(5)= 0.1,	XUNIF_ALBVIS_SOIL	(6)= 0.1,
XUNIF_ALBUV_SOIL	(4)= 0.06,	XUNIF_ALBUV_SOIL	(5)= 0.06,	XUNIF_ALBUV_SOIL	(6)= 0.06,
XUNIF_GMES	(4)= 0.003,	XUNIF_GMES	(5)= 0.003,	XUNIF_GMES	(6)= 0.003,
XUNIF_RE25	(4)= 0.0000003,	XUNIF_RE25	(5)= 0.0000003,	XUNIF_RE25	(6)= 0.0000003,
XUNIF_BSLAI	(4)= 0.06,	XUNIF_BSLAI	(5)= 0.06,	XUNIF_BSLAI	(6)= 0.06,
XUNIF_LAIMIN	(4)= 0.3,	XUNIF_LAIMIN	(5)= 0.3,	XUNIF_LAIMIN	(6)= 0.3,
XUNIF_SEFOLD	(4)= 5184000.,	XUNIF_SEFOLD	(5)= 5184000.,	XUNIF_SEFOLD	(6)= 5184000.,
XUNIF_GC	(4)= 0.00025,	XUNIF_GC	(5)= 0.00025,	XUNIF_GC	(6)= 0.00025,
XUNIF_DMAX	(4)= 0.1,	XUNIF_DMAX	(5)= 0.1,	XUNIF_DMAX	(6)= 0.1,
XUNIF_F2I	(4)= 0.3,	XUNIF_F2I	(5)= 0.3,	XUNIF_F2I	(6)= 0.3,
XUNIF_H_TREE	(4)= 20.,	XUNIF_H_TREE	(5)= 20.,	XUNIF_H_TREE	(6)= 20.,
XUNIF_CE_NITRO	(4)= 3.79,	XUNIF_CE_NITRO	(5)= 3.79,	XUNIF_CE_NITRO	(6)= 3.79,
XUNIF_CF_NITRO	(4)= 9.84,	XUNIF_CF_NITRO	(5)= 9.84,	XUNIF_CF_NITRO	(6)= 9.84,
XUNIF_CNA_NITRO	(4)= 1.3,	XUNIF_CNA_NITRO	(5)= 1.3,	XUNIF_CNA_NITRO	(6)= 1.3,
XUNIF_RSMIN	(7)= 40.,	XUNIF_RSMIN	(8)= 40.,	XUNIF_RSMIN	(9)= 40.,
XUNIF_GAMMA	(7)= 0.,	XUNIF_GAMMA	(8)= 0.,	XUNIF_GAMMA	(9)= 0.,
XUNIF_WRMAX_CF	(7)= 0.2,	XUNIF_WRMAX_CF	(8)= 0.2,	XUNIF_WRMAX_CF	(9)= 0.2,

XUNIF_RGL	(7)= 100.,	XUNIF_RGL	(8)= 100.,	XUNIF_RGL	(9)= 100.,
XUNIF_CV	(7)= 0.00002,	XUNIF_CV	(8)= 0.00002,	XUNIF_CV	(9)= 0.00002,
XUNIF_ZO_0_ZOH	(7)= 10.,	XUNIF_ZO_0_ZOH	(8)= 10.,	XUNIF_ZO_0_ZOH	(9)= 10.,
XUNIF_ALBNIR_VEG	(7)= 0.3,	XUNIF_ALBNIR_VEG	(8)= 0.3,	XUNIF_ALBNIR_VEG	(9)= 0.3,
XUNIF_ALBVIS_VEG	(7)= 0.1,	XUNIF_ALBVIS_VEG	(8)= 0.1,	XUNIF_ALBVIS_VEG	(9)= 0.1,
XUNIF_ALBUV_VEG	(7)= 0.0425,	XUNIF_ALBUV_VEG	(8)= 0.0425,	XUNIF_ALBUV_VEG	(9)= 0.0425,
XUNIF_ALBNIR_SOIL	(7)= 0.3,	XUNIF_ALBNIR_SOIL	(8)= 0.3,	XUNIF_ALBNIR_SOIL	(9)= 0.3,
XUNIF_ALBVIS_SOIL	(7)= 0.1,	XUNIF_ALBVIS_SOIL	(8)= 0.1,	XUNIF_ALBVIS_SOIL	(9)= 0.1,
XUNIF_ALBUV_SOIL	(7)= 0.06,	XUNIF_ALBUV_SOIL	(8)= 0.06,	XUNIF_ALBUV_SOIL	(9)= 0.06,
XUNIF_GMES	(7)= 0.003,	XUNIF_GMES	(8)= 0.003,	XUNIF_GMES	(9)= 0.003,
XUNIF_RE25	(7)= 0.0000003,	XUNIF_RE25	(8)= 0.0000003,	XUNIF_RE25	(9)= 0.0000003,
XUNIF_BSLAI	(7)= 0.06,	XUNIF_BSLAI	(8)= 0.06,	XUNIF_BSLAI	(9)= 0.06,
XUNIF_LAIMIN	(7)= 0.3,	XUNIF_LAIMIN	(8)= 0.3,	XUNIF_LAIMIN	(9)= 0.3,
XUNIF_SEFOLD	(7)= 5184000.,	XUNIF_SEFOLD	(8)= 5184000.,	XUNIF_SEFOLD	(9)= 5184000.,
XUNIF_GC	(7)= 0.00025,	XUNIF_GC	(8)= 0.00025,	XUNIF_GC	(9)= 0.00025,
XUNIF_DMAX	(7)= 0.1,	XUNIF_DMAX	(8)= 0.1,	XUNIF_DMAX	(9)= 0.1,
XUNIF_F2I	(7)= 0.3,	XUNIF_F2I	(8)= 0.3,	XUNIF_F2I	(9)= 0.3,
XUNIF_H_TREE	(7)= 20.,	XUNIF_H_TREE	(8)= 20.,	XUNIF_H_TREE	(9)= 20.,
XUNIF_CE_NITRO	(7)= 3.79,	XUNIF_CE_NITRO	(8)= 3.79,	XUNIF_CE_NITRO	(9)= 3.79,
XUNIF_CF_NITRO	(7)= 9.84,	XUNIF_CF_NITRO	(8)= 9.84,	XUNIF_CF_NITRO	(9)= 9.84,
XUNIF_CNA_NITRO	(7)= 1.3,	XUNIF_CNA_NITRO	(8)= 1.3,	XUNIF_CNA_NITRO	(9)= 1.3,
XUNIF_RSMIN	(10)= 40.,	XUNIF_RSMIN	(11)= 40.,	XUNIF_RSMIN	(12)= 40.,
XUNIF_GAMMA	(10)= 0.,	XUNIF_GAMMA	(11)= 0.,	XUNIF_GAMMA	(12)= 0.,
XUNIF_WRMAX_CF	(10)= 0.2,	XUNIF_WRMAX_CF	(11)= 0.2,	XUNIF_WRMAX_CF	(12)= 0.2,
XUNIF_RGL	(10)= 100.,	XUNIF_RGL	(11)= 100.,	XUNIF_RGL	(12)= 100.,
XUNIF_CV	(10)= 0.00002,	XUNIF_CV	(11)= 0.00002,	XUNIF_CV	(12)= 0.00002,
XUNIF_ZO_0_ZOH	(10)= 10.,	XUNIF_ZO_0_ZOH	(11)= 10.,	XUNIF_ZO_0_ZOH	(12)= 10.,
XUNIF_ALBNIR_VEG	(10)= 0.3,	XUNIF_ALBNIR_VEG	(11)= 0.3,	XUNIF_ALBNIR_VEG	(12)= 0.3,
XUNIF_ALBVIS_VEG	(10)= 0.1,	XUNIF_ALBVIS_VEG	(11)= 0.1,	XUNIF_ALBVIS_VEG	(12)= 0.1,
XUNIF_ALBUV_VEG	(10)= 0.0425,	XUNIF_ALBUV_VEG	(11)= 0.0425,	XUNIF_ALBUV_VEG	(12)= 0.0425,
XUNIF_ALBNIR_SOIL	(10)= 0.3,	XUNIF_ALBNIR_SOIL	(11)= 0.3,	XUNIF_ALBNIR_SOIL	(12)= 0.3,
XUNIF_ALBVIS_SOIL	(10)= 0.1,	XUNIF_ALBVIS_SOIL	(11)= 0.1,	XUNIF_ALBVIS_SOIL	(12)= 0.1,
XUNIF_ALBUV_SOIL	(10)= 0.06,	XUNIF_ALBUV_SOIL	(11)= 0.06,	XUNIF_ALBUV_SOIL	(12)= 0.06,
XUNIF_GMES	(10)= 0.003,	XUNIF_GMES	(11)= 0.003,	XUNIF_GMES	(12)= 0.003,
XUNIF_RE25	(10)= 0.0000003,	XUNIF_RE25	(11)= 0.0000003,	XUNIF_RE25	(12)= 0.0000003,
XUNIF_BSLAI	(10)= 0.06,	XUNIF_BSLAI	(11)= 0.06,	XUNIF_BSLAI	(12)= 0.06,
XUNIF_LAIMIN	(10)= 0.3,	XUNIF_LAIMIN	(11)= 0.3,	XUNIF_LAIMIN	(12)= 0.3,
XUNIF_SEFOLD	(10)= 5184000.,	XUNIF_SEFOLD	(11)= 5184000.,	XUNIF_SEFOLD	(12)= 5184000.,
XUNIF_GC	(10)= 0.00025,	XUNIF_GC	(11)= 0.00025,	XUNIF_GC	(12)= 0.00025,
XUNIF_DMAX	(10)= 0.1,	XUNIF_DMAX	(11)= 0.1,	XUNIF_DMAX	(12)= 0.1,
XUNIF_F2I	(10)= 0.3,	XUNIF_F2I	(11)= 0.3,	XUNIF_F2I	(12)= 0.3,
XUNIF_H_TREE	(10)= 20.,	XUNIF_H_TREE	(11)= 20.,	XUNIF_H_TREE	(12)= 20.,
XUNIF_CE_NITRO	(10)= 3.79,	XUNIF_CE_NITRO	(11)= 3.79,	XUNIF_CE_NITRO	(12)= 3.79,
XUNIF_CF_NITRO	(10)= 9.84,	XUNIF_CF_NITRO	(11)= 9.84,	XUNIF_CF_NITRO	(12)= 9.84,
XUNIF_CNA_NITRO	(10)= 1.3,	XUNIF_CNA_NITRO	(11)= 1.3,	XUNIF_CNA_NITRO	(12)= 1.3,

/
&NAM_DATA_TEB

```

NROOF_LAYER      = 3,
XUNIF_ALB_ROOF   = 0.2,
XUNIF_EMIS_ROOF  = 0.97,
XUNIF_HC_ROOF(1) = 2110000.,
XUNIF_HC_ROOF(2) = 2800000.,
XUNIF_HC_ROOF(3) = 2900000.,
XUNIF_TC_ROOF(1) = 1.51,
XUNIF_TC_ROOF(2) = 0.08,
XUNIF_TC_ROOF(3) = 0.05,
XUNIF_D_ROOF(1)  = 0.05,
XUNIF_D_ROOF(2)  = 0.4,
XUNIF_D_ROOF(3)  = 0.1,
NROAD_LAYER      = 3,
XUNIF_ALB_ROAD   = 0.2,
XUNIF_EMIS_ROAD  = 0.97,
XUNIF_HC_ROAD(1) = 2110000.,
XUNIF_HC_ROAD(2) = 2800000.,
XUNIF_HC_ROAD(3) = 2900000.,
XUNIF_TC_ROAD(1) = 1.51,
XUNIF_TC_ROAD(2) = 0.08,
XUNIF_TC_ROAD(3) = 0.05,
XUNIF_D_ROAD(1)  = 0.05,

```

```

XUNIF_D_ROAD(2) = 0.4,
XUNIF_D_ROAD(3) = 0.1,
NWALL_LAYER      = 3,
XUNIF_ALB_WALL   = 0.2,
XUNIF_EMIS_WALL  = 0.97,
XUNIF_HC_WALL(1) = 2110000.,
XUNIF_HC_WALL(2) = 2800000.,
XUNIF_HC_WALL(3) = 2900000.,
XUNIF_TC_WALL(1) = 1.51,
XUNIF_TC_WALL(2) = 0.08,
XUNIF_TC_WALL(3) = 0.05,
XUNIF_D_WALL(1)  = 0.05,
XUNIF_D_WALL(2)  = 0.4,
XUNIF_D_WALL(3)  = 0.1,
XUNIF_ZO_TOWN    = 1.,
XUNIF_BLD        = 0.5,
XUNIF_BLD_HEIGHT = 10.,
XUNIF_WALL_O_HOR = 0.5,
XUNIF_H_TRAFFIC  = 10.,
XUNIF_LE_TRAFFIC = 0.,
XUNIF_H_INDUSTRY = 5.,
XUNIF_LE_INDUSTRY= 0.

/
&NAM_FRAC        LECOCLIMAP = F,
XUNIF_SEA        = 0.,
XUNIF_WATER      = 0.,
XUNIF_TOWN       = 0.,
XUNIF_NATURE     = 1.

/
&NAM_PGD_GRID    CGRID = 'LONLAT REG'

/
&NAM_LONLAT_REG  XLONMIN = 0.,           ,
XLONMAX = 0.,           ,
XLATMIN = 0.,           ,
XLATMAX = 0.,           ,
NLON      = 1           ,
NLAT      = 1

/
&NAM_PGD_SCHEMES CNATURE = 'ISBA  ',    ,
CSEA      = 'SEAFIX',    ,
CTOWN     = 'TEB  ',    ,
CWATER    = 'WATFLX'

/
&NAM_ZS          XUNIF_ZS = 113.

/
&NAM_ISBA        XUNIF_CLAY = 0.37      ,
XUNIF_SAND = 0.37      ,
XUNIF_RUNOFFB = 0.5    ,
CISBA        = '2-L'   ,
CPHOTO       = 'NIT'   ,
NPATCH       = 12      ,
NGROUND_LAYER = 2

/
&NAM_PREPFILE    CPREPFILE = 'PREP'

/
&NAM_PREP_SURF_ATM NYEAR = 1986,
NMONTH = 1,
NDAY   = 1,
XTIME  = 0.

/
&NAM_PREP_SEAFLUX XSST_UNIF = 285.,
NYEAR = 1986,
NMONTH = 1,
NDAY   = 1,
XTIME  = 0.

/
&NAM_PREP_WATFLUX XTS_WATER_UNIF = 285.,

```

```

        NYEAR = 1986,
        NMONTH = 1,
        NDAY = 1,
        XTIME = 0.
/
&NAM_PREP_TEB      XTI_ROAD= 285.,
                   XTI_BLD = 285.
                   XTS_ROAD= 285.
                   XTS_ROOF= 285.,
                   XTS_WALL= 285.,
                   XWS_ROAD= 0.,
                   XWS_ROOF= 0.,
                   NYEAR = 1986,
                   NMONTH = 1,
                   NDAY = 1,
                   XTIME = 0.
/
&NAM_PREP_ISBA     XHUG_SURF = 1.,
                   XHUG_ROOT = 1.,
                   XHUG_DEEP = 1.,
                   XTG_SURF = 276.16,
                   XTG_ROOT = 276.16,
                   XTG_DEEP = 276.16,
                   NYEAR = 1986,
                   NMONTH = 1,
                   NDAY = 1,
                   XTIME = 0.
/
&NAM_PREP_ISBA_SNOW CSNOW = '3-L'
/

&NAM_IO_OFFLINE    YPROGRAM = 'ASCII' ,
                   YCOUPLING = 'E'
/
&NAM_DIAG_SURFn    LSURF_BUDGET = F      ,
                   N2M          = 0      ,
                   LCOEF        = F      ,
                   LSURF_VARS   = F
/
&NAM_DIAG_SURF_ATMn LFRAC          = F
/
&NAM_DIAG_ISBAn    LPGD              = F ,
                   LSURF_EVAP_BUDGET = F ,
                   LSURF_MISC_BUDGET = F ,
                   LSURF_BUDGETC     = F
/
&NAM_DIAG_TEBn    LSURF_MISC_BUDGET = F
/
&NAM_SGH_ISBAn     CRUNOFF           = "WSAT"
/
&NAM_ISBAn         CROUGH            = "Z04D" ,
                   CSCOND            = "NP89" ,
                   CALBEDO           = "DRY" ,
                   CC1DRY            = 'DEF' ,
                   CSOILFRZ          = 'DEF' ,
                   CDIFSFCND         = 'DEF' ,
                   CSNOWRES          = 'DEF' ,
                   CCPSURF           = 'DRY'
/
&NAM_CH_ISBAn      CCH_DRY_DEP = "WES89 "
/
&NAM_SEAFLUXn     CSEA_ALB = "TA96"
/

```

```

&NAM_CH_SEAFLUXn      CCH_DRY_DEP = "WES89 "
/
&NAM_CH_WATFLUXn      CCH_DRY_DEP = "WES89 "
/
&NAM_CH_TEBn          CCH_DRY_DEP = "WES89 "
/

```

A.2.3 Surface parameters read from external files

The following namelist is valid only for simulation without patches. In case of use of patches (like for A-gs options), it should be updated.

```

&NAM_DATA_ISBA      NTIME = 12 ,
CFNAM_VEGTYPE(1)    = 'VEGTYPE_01.DAT', CFTYP_VEGTYPE(1)    = 'ASCLLV',
CFNAM_VEGTYPE(2)    = 'VEGTYPE_02.DAT', CFTYP_VEGTYPE(2)    = 'ASCLLV',
CFNAM_VEGTYPE(3)    = 'VEGTYPE_03.DAT', CFTYP_VEGTYPE(3)    = 'ASCLLV',
CFNAM_VEGTYPE(4)    = 'VEGTYPE_04.DAT', CFTYP_VEGTYPE(4)    = 'ASCLLV',
CFNAM_VEGTYPE(5)    = 'VEGTYPE_05.DAT', CFTYP_VEGTYPE(5)    = 'ASCLLV',
CFNAM_VEGTYPE(6)    = 'VEGTYPE_06.DAT', CFTYP_VEGTYPE(6)    = 'ASCLLV',
CFNAM_VEGTYPE(7)    = 'VEGTYPE_07.DAT', CFTYP_VEGTYPE(7)    = 'ASCLLV',
CFNAM_VEGTYPE(8)    = 'VEGTYPE_08.DAT', CFTYP_VEGTYPE(8)    = 'ASCLLV',
CFNAM_VEGTYPE(9)    = 'VEGTYPE_09.DAT', CFTYP_VEGTYPE(9)    = 'ASCLLV',
CFNAM_VEGTYPE(10)   = 'VEGTYPE_10.DAT', CFTYP_VEGTYPE(10)   = 'ASCLLV',
CFNAM_VEGTYPE(11)   = 'VEGTYPE_11.DAT', CFTYP_VEGTYPE(11)   = 'ASCLLV',
CFNAM_VEGTYPE(12)   = 'VEGTYPE_12.DAT', CFTYP_VEGTYPE(12)   = 'ASCLLV',
CFNAM_VEG(1,1)      = 'VEG_01.DAT', CFTYP_VEG(1,1)      = 'ASCLLV',
CFNAM_VEG(1,2)      = 'VEG_02.DAT', CFTYP_VEG(1,2)      = 'ASCLLV',
CFNAM_VEG(1,3)      = 'VEG_03.DAT', CFTYP_VEG(1,3)      = 'ASCLLV',
CFNAM_VEG(1,4)      = 'VEG_04.DAT', CFTYP_VEG(1,4)      = 'ASCLLV',
CFNAM_VEG(1,5)      = 'VEG_05.DAT', CFTYP_VEG(1,5)      = 'ASCLLV',
CFNAM_VEG(1,6)      = 'VEG_06.DAT', CFTYP_VEG(1,6)      = 'ASCLLV',
CFNAM_VEG(1,7)      = 'VEG_07.DAT', CFTYP_VEG(1,7)      = 'ASCLLV',
CFNAM_VEG(1,8)      = 'VEG_08.DAT', CFTYP_VEG(1,8)      = 'ASCLLV',
CFNAM_VEG(1,9)      = 'VEG_09.DAT', CFTYP_VEG(1,9)      = 'ASCLLV',
CFNAM_VEG(1,10)     = 'VEG_10.DAT', CFTYP_VEG(1,10)     = 'ASCLLV',
CFNAM_VEG(1,11)     = 'VEG_11.DAT', CFTYP_VEG(1,11)     = 'ASCLLV',
CFNAM_VEG(1,12)     = 'VEG_12.DAT', CFTYP_VEG(1,12)     = 'ASCLLV',
CFNAM_LAI(1,1)      = 'LAI_01.DAT', CFTYP_LAI(1,1)      = 'ASCLLV',
CFNAM_LAI(1,2)      = 'LAI_02.DAT', CFTYP_LAI(1,2)      = 'ASCLLV',
CFNAM_LAI(1,3)      = 'LAI_03.DAT', CFTYP_LAI(1,3)      = 'ASCLLV',
CFNAM_LAI(1,4)      = 'LAI_04.DAT', CFTYP_LAI(1,4)      = 'ASCLLV',
CFNAM_LAI(1,5)      = 'LAI_05.DAT', CFTYP_LAI(1,5)      = 'ASCLLV',
CFNAM_LAI(1,6)      = 'LAI_06.DAT', CFTYP_LAI(1,6)      = 'ASCLLV',
CFNAM_LAI(1,7)      = 'LAI_07.DAT', CFTYP_LAI(1,7)      = 'ASCLLV',
CFNAM_LAI(1,8)      = 'LAI_08.DAT', CFTYP_LAI(1,8)      = 'ASCLLV',
CFNAM_LAI(1,9)      = 'LAI_09.DAT', CFTYP_LAI(1,9)      = 'ASCLLV',
CFNAM_LAI(1,10)     = 'LAI_10.DAT', CFTYP_LAI(1,10)     = 'ASCLLV',
CFNAM_LAI(1,11)     = 'LAI_11.DAT', CFTYP_LAI(1,11)     = 'ASCLLV',
CFNAM_LAI(1,12)     = 'LAI_12.DAT', CFTYP_LAI(1,12)     = 'ASCLLV',
CFNAM_ZO(1,1)       = 'ZO_01.DAT', CFTYP_ZO(1,1)       = 'ASCLLV',
CFNAM_ZO(1,2)       = 'ZO_02.DAT', CFTYP_ZO(1,2)       = 'ASCLLV',
CFNAM_ZO(1,3)       = 'ZO_03.DAT', CFTYP_ZO(1,3)       = 'ASCLLV',
CFNAM_ZO(1,4)       = 'ZO_04.DAT', CFTYP_ZO(1,4)       = 'ASCLLV',
CFNAM_ZO(1,5)       = 'ZO_05.DAT', CFTYP_ZO(1,5)       = 'ASCLLV',
CFNAM_ZO(1,6)       = 'ZO_06.DAT', CFTYP_ZO(1,6)       = 'ASCLLV',
CFNAM_ZO(1,7)       = 'ZO_07.DAT', CFTYP_ZO(1,7)       = 'ASCLLV',
CFNAM_ZO(1,8)       = 'ZO_08.DAT', CFTYP_ZO(1,8)       = 'ASCLLV',
CFNAM_ZO(1,9)       = 'ZO_09.DAT', CFTYP_ZO(1,9)       = 'ASCLLV',
CFNAM_ZO(1,10)      = 'ZO_10.DAT', CFTYP_ZO(1,10)      = 'ASCLLV',
CFNAM_ZO(1,11)      = 'ZO_11.DAT', CFTYP_ZO(1,11)      = 'ASCLLV',
CFNAM_ZO(1,12)      = 'ZO_12.DAT', CFTYP_ZO(1,12)      = 'ASCLLV',
CFNAM_EMIS(1,1)     = 'EMIS_01.DAT', CFTYP_EMIS(1,1)     = 'ASCLLV',
CFNAM_EMIS(1,2)     = 'EMIS_02.DAT', CFTYP_EMIS(1,2)     = 'ASCLLV',
CFNAM_EMIS(1,3)     = 'EMIS_03.DAT', CFTYP_EMIS(1,3)     = 'ASCLLV',
CFNAM_EMIS(1,4)     = 'EMIS_04.DAT', CFTYP_EMIS(1,4)     = 'ASCLLV',
CFNAM_EMIS(1,5)     = 'EMIS_05.DAT', CFTYP_EMIS(1,5)     = 'ASCLLV',
CFNAM_EMIS(1,6)     = 'EMIS_06.DAT', CFTYP_EMIS(1,6)     = 'ASCLLV',
CFNAM_EMIS(1,7)     = 'EMIS_07.DAT', CFTYP_EMIS(1,7)     = 'ASCLLV',
CFNAM_EMIS(1,8)     = 'EMIS_08.DAT', CFTYP_EMIS(1,8)     = 'ASCLLV',

```

CFNAM_EMIS(1,9)	=	'EMIS_09.DAT	,	CFTYP_EMIS(1,9)	=	'ASCLLV'
CFNAM_EMIS(1,10)	=	'EMIS_10.DAT	,	CFTYP_EMIS(1,10)	=	'ASCLLV'
CFNAM_EMIS(1,11)	=	'EMIS_11.DAT	,	CFTYP_EMIS(1,11)	=	'ASCLLV'
CFNAM_EMIS(1,12)	=	'EMIS_12.DAT	,	CFTYP_EMIS(1,12)	=	'ASCLLV'
CFNAM_DG(1,1)	=	'DG_1.DAT	,	CFTYP_DG(1,1)	=	'ASCLLV'
CFNAM_DG(1,2)	=	'DG_2.DAT	,	CFTYP_DG(1,2)	=	'ASCLLV'
CFNAM_DG(1,3)	=	'DG_3.DAT	,	CFTYP_DG(1,3)	=	'ASCLLV'
CFNAM_ROOTFRAC(1,1)	=	'ROOTFRAC_1.DAT	,	CFTYP_ROOTFRAC(1,1)	=	'ASCLLV'
CFNAM_ROOTFRAC(1,2)	=	'ROOTFRAC_2.DAT	,	CFTYP_ROOTFRAC(1,2)	=	'ASCLLV'
CFNAM_ROOTFRAC(1,3)	=	'ROOTFRAC_3.DAT	,	CFTYP_ROOTFRAC(1,3)	=	'ASCLLV'
CFNAM_RSMIN(1)	=	'RSMIN.DAT	,	CFTYP_RSMIN(1)	=	'ASCLLV'
CFNAM_ZO_0_ZOH(1)	=	'ZO_0_ZOH.DAT	,	CFTYP_ZO_0_ZOH(1)	=	'ASCLLV'
CFNAM_GAMMA(1)	=	'GAMMA.DAT	,	CFTYP_GAMMA(1)	=	'ASCLLV'
CFNAM_WRMX_CF(1)	=	'WRMX_CF.DAT	,	CFTYP_WRMX_CF(1)	=	'ASCLLV'
CFNAM_RGL(1)	=	'RGL.DAT	,	CFTYP_RGL(1)	=	'ASCLLV'
CFNAM_CV(1)	=	'CV.DAT	,	CFTYP_CV(1)	=	'ASCLLV'
CFNAM_ALBNIR_VEG(1)	=	'ALBNIR_VEG.DAT	,	CFTYP_ALBNIR_VEG(1)	=	'ASCLLV'
CFNAM_ALBVIS_VEG(1)	=	'ALBVIS_VEG.DAT	,	CFTYP_ALBVIS_VEG(1)	=	'ASCLLV'
CFNAM_ALBUV_VEG(1)	=	'ALBUV_VEG.DAT	,	CFTYP_ALBUV_VEG(1)	=	'ASCLLV'
CFNAM_ALBNIR_SOIL(1)	=	'ALBNIR_SOIL.DAT	,	CFTYP_ALBNIR_SOIL(1)	=	'ASCLLV'
CFNAM_ALBVIS_SOIL(1)	=	'ALBVIS_SOIL.DAT	,	CFTYP_ALBVIS_SOIL(1)	=	'ASCLLV'
CFNAM_ALBUV_SOIL(1)	=	'ALBUV_SOIL.DAT	,	CFTYP_ALBUV_SOIL(1)	=	'ASCLLV'
CFNAM_GMES(1)	=	'GMES.DAT	,	CFTYP_GMES(1)	=	'ASCLLV'
CFNAM_RE25(1)	=	'RE25.DAT	,	CFTYP_RE25(1)	=	'ASCLLV'
CFNAM_BSLAI(1)	=	'BSLAI.DAT	,	CFTYP_BSLAI(1)	=	'ASCLLV'
CFNAM_LAIMIN(1)	=	'LAIMIN.DAT	,	CFTYP_LAIMIN(1)	=	'ASCLLV'
CFNAM_SEFOLD(1)	=	'SEFOLD.DAT	,	CFTYP_SEFOLD(1)	=	'ASCLLV'
CFNAM_GC(1)	=	'GC.DAT	,	CFTYP_GC(1)	=	'ASCLLV'
CFNAM_DMAX(1)	=	'DMAX.DAT	,	CFTYP_DMAX(1)	=	'ASCLLV'
CFNAM_F2I(1)	=	'F2I.DAT	,	CFTYP_F2I(1)	=	'ASCLLV'
CFNAM_H_TREE(1)	=	'H_TREE.DAT	,	CFTYP_H_TREE(1)	=	'ASCLLV'
CFNAM_CE_NITRO(1)	=	'CE_NITRO.DAT	,	CFTYP_CE_NITRO(1)	=	'ASCLLV'
CFNAM_CF_NITRO(1)	=	'CF_NITRO.DAT	,	CFTYP_CF_NITRO(1)	=	'ASCLLV'
CFNAM_CNA_NITRO(1)	=	'CNA_NITRO.DAT	,	CFTYP_CNA_NITRO(1)	=	'ASCLLV'
/						
&NAM_DATA_TEB	NROOF_LAYER = 3 ,					
	CFNAM_ALB_ROOF	= 'ALB_ROOF.DAT	,	CFTYP_ALB_ROOF	= 'ASCLLV'	
	CFNAM_EMIS_ROOF	= 'EMIS_ROOF.DAT	,	CFTYP_EMIS_ROOF	= 'ASCLLV'	
	CFNAM_HC_ROOF(1)	= 'HC_ROOF.DAT	,	CFTYP_HC_ROOF(1)	= 'ASCLLV'	
	CFNAM_HC_ROOF(2)	= 'HC_ROOF.DAT	,	CFTYP_HC_ROOF(2)	= 'ASCLLV'	
	CFNAM_HC_ROOF(3)	= 'HC_ROOF.DAT	,	CFTYP_HC_ROOF(3)	= 'ASCLLV'	
	CFNAM_TC_ROOF(1)	= 'TC_ROOF.DAT	,	CFTYP_TC_ROOF(1)	= 'ASCLLV'	
	CFNAM_TC_ROOF(2)	= 'TC_ROOF.DAT	,	CFTYP_TC_ROOF(2)	= 'ASCLLV'	
	CFNAM_TC_ROOF(3)	= 'TC_ROOF.DAT	,	CFTYP_TC_ROOF(3)	= 'ASCLLV'	
	CFNAM_D_ROOF(1)	= 'D_ROOF.DAT	,	CFTYP_D_ROOF(1)	= 'ASCLLV'	
	CFNAM_D_ROOF(2)	= 'D_ROOF.DAT	,	CFTYP_D_ROOF(2)	= 'ASCLLV'	
	CFNAM_D_ROOF(3)	= 'D_ROOF.DAT	,	CFTYP_D_ROOF(3)	= 'ASCLLV'	
	NROAD_LAYER = 3 ,					
	CFNAM_ALB_ROAD	= 'ALB_ROAD.DAT	,	CFTYP_ALB_ROAD	= 'ASCLLV'	
	CFNAM_EMIS_ROAD	= 'EMIS_ROAD.DAT	,	CFTYP_EMIS_ROAD	= 'ASCLLV'	
	CFNAM_HC_ROAD(1)	= 'HC_ROAD.DAT	,	CFTYP_HC_ROAD(1)	= 'ASCLLV'	
	CFNAM_HC_ROAD(2)	= 'HC_ROAD.DAT	,	CFTYP_HC_ROAD(2)	= 'ASCLLV'	
	CFNAM_HC_ROAD(3)	= 'HC_ROAD.DAT	,	CFTYP_HC_ROAD(3)	= 'ASCLLV'	
	CFNAM_TC_ROAD(1)	= 'TC_ROAD.DAT	,	CFTYP_TC_ROAD(1)	= 'ASCLLV'	
	CFNAM_TC_ROAD(2)	= 'TC_ROAD.DAT	,	CFTYP_TC_ROAD(2)	= 'ASCLLV'	
	CFNAM_TC_ROAD(3)	= 'TC_ROAD.DAT	,	CFTYP_TC_ROAD(3)	= 'ASCLLV'	
	CFNAM_D_ROAD(1)	= 'D_ROAD.DAT	,	CFTYP_D_ROAD(1)	= 'ASCLLV'	
	CFNAM_D_ROAD(2)	= 'D_ROAD.DAT	,	CFTYP_D_ROAD(2)	= 'ASCLLV'	
	CFNAM_D_ROAD(3)	= 'D_ROAD.DAT	,	CFTYP_D_ROAD(3)	= 'ASCLLV'	
	NWALL_LAYER = 3 ,					
	CFNAM_ALB_WALL	= 'ALB_WALL.DAT	,	CFTYP_ALB_WALL	= 'ASCLLV'	
	CFNAM_EMIS_WALL	= 'EMIS_WALL.DAT	,	CFTYP_EMIS_WALL	= 'ASCLLV'	
	CFNAM_HC_WALL(1)	= 'HC_WALL.DAT	,	CFTYP_HC_WALL(1)	= 'ASCLLV'	
	CFNAM_HC_WALL(2)	= 'HC_WALL.DAT	,	CFTYP_HC_WALL(2)	= 'ASCLLV'	
	CFNAM_HC_WALL(3)	= 'HC_WALL.DAT	,	CFTYP_HC_WALL(3)	= 'ASCLLV'	
	CFNAM_TC_WALL(1)	= 'TC_WALL.DAT	,	CFTYP_TC_WALL(1)	= 'ASCLLV'	
	CFNAM_TC_WALL(2)	= 'TC_WALL.DAT	,	CFTYP_TC_WALL(2)	= 'ASCLLV'	
	CFNAM_TC_WALL(3)	= 'TC_WALL.DAT	,	CFTYP_TC_WALL(3)	= 'ASCLLV'	

```

CFNAM_D_WALL(1)      = 'D_WALL.DAT'          , CFTYP_D_WALL(1)      = 'ASCLLV',
CFNAM_D_WALL(2)      = 'D_WALL.DAT'          , CFTYP_D_WALL(2)      = 'ASCLLV',
CFNAM_D_WALL(3)      = 'D_WALL.DAT'          , CFTYP_D_WALL(3)      = 'ASCLLV',
CFNAM_ZO_TOWN         = 'ZO_TOWN.DAT'         , CFTYP_ZO_TOWN        = 'ASCLLV',
CFNAM_BLD             = 'BLD.DAT'             , CFTYP_BLD            = 'ASCLLV',
CFNAM_BLD_HEIGHT      = 'BLD_HEIGHT.DAT'      , CFTYP_BLD_HEIGHT     = 'ASCLLV',
CFNAM_WALL_O_HOR      = 'WALL_O_HOR.DAT'      , CFTYP_WALL_O_HOR     = 'ASCLLV',
CFNAM_H_TRAFFIC       = 'H_TRAFFIC.DAT'       , CFTYP_H_TRAFFIC      = 'ASCLLV',
CFNAM_LE_TRAFFIC      = 'LE_TRAFFIC.DAT'      , CFTYP_LE_TRAFFIC     = 'ASCLLV',
CFNAM_H_INDUSTRY      = 'H_INDUSTRY.DAT'      , CFTYP_H_INDUSTRY     = 'ASCLLV',
CFNAM_LE_INDUSTRY     = 'LE_INDUSTRY.DAT'     , CFTYP_LE_INDUSTRY    = 'ASCLLV',

/
&NAM_FRAC             LECOCLIMAP = F,
CFNAM_SEA              = 'SEA.DAT'            , CFTYP_SEA            = 'ASCLLV',
CFNAM_WATER            = 'WATER.DAT'          , CFTYP_WATER          = 'ASCLLV',
CFNAM_NATURE           = 'NATURE.DAT'         , CFTYP_NATURE         = 'ASCLLV',
CFNAM_TOWN             = 'TOWN.DAT'           , CFTYP_TOWN           = 'ASCLLV',

/
&NAM_PGD_GRID         CGRID = 'LONLAT REG'

/
&NAM_LONLAT_REG       XLONMIN = 0.           ,
                        XLONMAX = 0.5         ,
                        XLATMIN = 0.           ,
                        XLATMAX = 0.5         ,
                        NLON   = 1             ,
                        NLAT   = 1

/
&NAM_PGD_SCHEMES      CNATURE = 'ISBA'      ,
                        CSEA    = 'SEAFLEX'   ,
                        CTOWN   = 'TEB'      ,
                        CWATER  = 'WATFLX'

/
&NAM_ZS               XUNIF_ZS = 0.

/
&NAM_ISBA             XUNIF_CLAY = 0.4        ,
                        XUNIF_SAND = 0.2        ,
                        XUNIF_RUNOFFB = 0.5      ,
                        CISBA     = '3-L'      ,
                        CPHOTO    = 'NON'      ,
                        NPATCH   = 1          ,
                        NGROUND_LAYER = 3

/
&NAM_PREPFILE         CPREPFILE = 'PREP'

/
&NAM_PREP_SURF_ATM    NYEAR = 2004,
                        NMONTH = 10,
                        NDAY   = 25,
                        XTIME  = 21600.

/
&NAM_PREP_SEAFLUX     XSST_UNIF = 285.,
                        NYEAR = 2004,
                        NMONTH = 10,
                        NDAY   = 25,
                        XTIME  = 21600.

/
&NAM_PREP_WATFLUX     XTS_WATER_UNIF = 285.,
                        NYEAR = 2004,
                        NMONTH = 10,
                        NDAY   = 25,
                        XTIME  = 21600.

/
&NAM_PREP_TEB         XTI_ROAD= 285.,
                        XTI_BLD = 285.,
                        XTS_ROAD= 285.,
                        XTS_ROOF= 285.,
                        XTS_WALL= 285.,

```

```

XWS_ROAD= 0.,
XWS_ROOF= 0.,
NYEAR  = 2004,
NMONTH = 10,
NDAY   = 25,
XTIME  = 21600.

/
&NAM_PREP_ISBA    XHUG_SURF = 0.2,
                  XHUG_ROOT = 0.2,
                  XHUG_DEEP = 0.2,
                  XTG_SURF  = 285.,
                  XTG_ROOT  = 288.,
                  XTG_DEEP  = 292.
                  NYEAR    = 2004,
                  NMONTH   = 10,
                  NDAY     = 25,
                  XTIME    = 21600.

/
&NAM_PREP_ISBA_SNOW    CSNOW = '3-L'

/


&NAM_IO_OFFLINE        YPROGRAM = 'ASCII' ,
                        YCOUPLING = 'E'

/
&NAM_DIAG_SURFh        LSURF_BUDGET = T      ,
                        N2M          = 1

/
&NAM_DIAG_SURF_ATMh    LFRAC          = T

/
&NAM_DIAG_ISBAh        LPGD            = T ,
                        LSURF_EVAP_BUDGET = T ,
                        LSURF_MISC_BUDGET = T ,
                        LSURF_BUDGETC    = F

/
&NAM_DIAG_TEBh         LSURF_MISC_BUDGET = T

/
&NAM_SGH_ISBAh         CRUNOFF         = "WSAT"

/
&NAM_ISBAh             CROUGH          = "Z04D" ,
                        CSCOND          = "NP89" ,
                        CALBEDO         = "DRY" ,
                        CC1DRY          = 'DEF' ,
                        CSOILFRZ        = 'DEF' ,
                        CDIFSFCND       = 'DEF' ,
                        CSNOWRES         = 'DEF' ,
                        CCPSURF         = 'DRY'

/
&NAM_CH_ISBAh          CCH_DRY_DEP     = "WES89 "

/
&NAM_SEAFLUXh          CSEA_ALB        = "TA96"

/
&NAM_CH_SEAFLUXh       CCH_DRY_DEP     = "WES89 "

/
&NAM_CH_WATFLUXh       CCH_DRY_DEP     = "WES89 "

/
&NAM_CH_TEBh           CCH_DRY_DEP     = "WES89 "

/

```

Index

- namelist description		NAM_CH_TEBn	72
NAM_CARTESIAN	49	NAM_CH_WATFLUXn	71
NAM_CH_CONTROLn	71	CCHEM_SURF_FILE	
NAM_CH_EMIS_PGD	57	NAM_CH_SURFn	71
NAM_CH_ISBAn	72	CCPSURF	
NAM_CH_SEAFLUXn	71	NAM_ISBAn	70
NAM_CH_SURFn	71	CDIFSFCND	
NAM_CH_TEBn	72	NAM_ISBAn	70
NAM_CH_WATFLUXn	71	CDUMMY_PGD_AREA	
NAM_CONF_PROJ	48	NAM_DUMMY_PGD	56
NAM_CONF_PROJ_GRID	48	CDUMMY_PGD_ATYPE	
NAM_COVER	53	NAM_DUMMY_PGD	56
NAM_DATA_FLAKE	45	CDUMMY_PGD_FILE	
NAM_DATA_ISBA	38	NAM_DUMMY_PGD	56
NAM_DATA_SEAFLUX	44	CDUMMY_PGD_FILETYPE	
NAM_DATA_TEB	42	NAM_DUMMY_PGD	56
NAM_DIAG_ISBAn	75	CDUMMY_PGD_NAME	
NAM_DIAG_SURF_ATMn	73	NAM_DUMMY_PGD	56
NAM_DIAG_SURFn	74	CEMIS_PGD_AREA	
NAM_DIAG_TEBn	75	NAM_CH_EMIS_PGD	57
NAM_DUMMY_PGD	56	CEMIS_PGD_ATYPE	
NAM_FLAKEn	69	NAM_CH_EMIS_PGD	57
NAM_FRAC	37	CEMIS_PGD_COMMENT	
NAM_IGN	51	NAM_CH_EMIS_PGD	57
NAM_INIFILE_CARTESIAN	49	CEMIS_PGD_FILE	
NAM_INIFILE_CONF_PROJ	48	NAM_CH_EMIS_PGD	57
NAM_ISBA	55	CEMIS_PGD_FILETYPE	
NAM_ISBAn	70	NAM_CH_EMIS_PGD	57
NAM_LONLAT_REG	50	CEMIS_PGD_NAME	
NAM_PGD_GRID	47	NAM_CH_EMIS_PGD	57
NAM_PGD_SCHEMES	46	CEMISPARAM	
NAM_PGDFILE	36	NAM_SURF_DST	70
NAM_PREP_FLAKE	63	CFILE	
NAM_PREP_ISBA	64	NAM_PREP_SURF_ATM	60
NAM_PREP_ISBA_SNOW	64	CFILE_FLAKE	
NAM_PREP_SEAFLUX	61	NAM_PREP_FLAKE	63
NAM_PREP_SURF_ATM	60	CFILE_ISBA	
NAM_PREP_TEB	65	NAM_PREP_ISBA	64
NAM_PREP_WATFLUX	62	CFILE_SEAFLX	
NAM_SEAFLUXn	68	NAM_PREP_SEAFLX	61
NAM_SGH_ISBAn	70	CFILE_TEB	
NAM_SURF_ATM	67	NAM_PREP_TEB	65
NAM_SURF_DST	70	CFILE_TG	
NAM_ZS	54	NAM_PREP_ISBA	64
NAMDIM	52	CFILE_TS	
NAMGEM	52	NAM_PREP_TEB	65
NAMRGRI	52	CFILE_WATFLX	
		NAM_PREP_WATFLX	62
		CFILE_WG	
		NAM_PREP_ISBA	64
		CFILE_WS	
		NAM_PREP_TEB	65
		CFILETYPE	
		NAM_PREP_SURF_ATM	60
		CFLAKE_FLUX	
		NAM_FLAKEn	69
		CFLAKE_SNOW	
C			
CALBEDO			
NAM_ISBAn	70		
CC1DRY			
NAM_ISBAn	70		
CCH_DRY_DEP			
NAM_CH_ISBAn	72		
NAM_CH_SEAFLUXn	71		

NAM_FLAKEn	69	NAM_DATA_TEB	42
CFNAM_		CFNAM_HC_ROOF	42
NAM_FRAC	37	NAM_DATA_TEB	42
CFNAM_ALB_ROAD		CFNAM_HC_WALL	43
NAM_DATA_TEB	42	NAM_DATA_TEB	43
CFNAM_ALB_ROOF		CFNAM_LAI	39
NAM_DATA_ISBA	42	NAM_DATA_ISBA	39
CFNAM_ALB_WALL		CFNAM_LAIMIN	41
NAM_DATA_TEB	43	NAM_DATA_ISBA	41
CFNAM_ALBNIR_SOIL		CFNAM_LE_INDUSTRIES	43
NAM_DATA_ISBA	40	NAM_DATA_TEB	43
CFNAM_ALBNIR_VEG		CFNAM_LE_TRAFFIC	43
NAM_DATA_ISBA	40	NAM_DATA_TEB	43
CFNAM_ALBUV_SOIL		CFNAM_RE25	41
NAM_DATA_ISBA	40	NAM_DATA_ISBA	41
CFNAM_ALBUV_VEG		CFNAM_RGL	40
NAM_DATA_ISBA	40	NAM_DATA_ISBA	40
CFNAM_ALBVIS_SOIL		CFNAM_ROOTFRAC	39
NAM_DATA_ISBA	40	NAM_DATA_ISBA	39
CFNAM_ALBVIS_VEG		CFNAM_RSMIN	40
NAM_DATA_ISBA	40	NAM_DATA_ISBA	40
CFNAM_BLD		CFNAM_SEA	37
NAM_DATA_TEB	42	NAM_FRAC	37
CFNAM_BLD_HEIGHT		CFNAM_SEFOLD	41
NAM_DATA_TEB	42	NAM_DATA_ISBA	41
CFNAM_BSLAI		CFNAM_SST	44
NAM_DATA_ISBA	41	NAM_DATA_SEAFLUX	44
CFNAM_CE_NITRO		CFNAM_TC_ROAD	42
NAM_DATA_ISBA	41	NAM_DATA_TEB	42
CFNAM_CF_NITRO		CFNAM_TC_ROOF	42
NAM_DATA_ISBA	41	NAM_DATA_TEB	43
CFNAM_CNA_NITRO		CFNAM_TC_WALL	39
NAM_DATA_ISBA	41	NAM_DATA_ISBA	39
CFNAM_CV		CFNAM_VEG	38
NAM_DATA_ISBA	40	NAM_DATA_ISBA	38
CFNAM_D_ROAD		CFNAM_VEGTYPE	42
NAM_DATA_TEB	42	NAM_DATA_ISBA	42
CFNAM_D_ROOF		CFNAM_WALL_O_HOR	42
NAM_DATA_TEB	42	NAM_DATA_TEB	42
CFNAM_D_WALL		CFNAM_WRMALX_CF	40
NAM_DATA_TEB	43	NAM_DATA_ISBA	40
CFNAM_DG		CFNAM_Z0	39
NAM_DATA_ISBA	39	NAM_DATA_ISBA	39
CFNAM_DMAX		CFNAM_Z0_O_ZOH	40
NAM_DATA_ISBA	41	NAM_DATA_ISBA	40
CFNAM_EMIS		CFNAM_Z0_TOWN	42
NAM_DATA_ISBA	39	NAM_DATA_TEB	42
CFNAM_EMIS_ROAD		CFTYP_ALB_ROAD	42
NAM_DATA_TEB	42	NAM_DATA_TEB	42
CFNAM_EMIS_ROOF		CFTYP_ALB_ROOF	42
NAM_DATA_TEB	42	NAM_DATA_TEB	43
CFNAM_EMIS_WALL		CFTYP_ALB_WALL	43
NAM_DATA_TEB	43	NAM_DATA_TEB	40
CFNAM_F2I		CFTYP_ALBNIR_SOIL	40
NAM_DATA_ISBA	41	NAM_DATA_ISBA	40
CFNAM_GAMMA		CFTYP_ALBNIR_VEG	40
NAM_DATA_ISBA	40	NAM_DATA_ISBA	40
CFNAM_GC		CFTYP_ALBUV_SOIL	40
NAM_DATA_ISBA	41	NAM_DATA_ISBA	40
CFNAM_GMES		CFTYP_ALBUV_VEG	40
NAM_DATA_ISBA	41	NAM_DATA_ISBA	40
CFNAM_H_INDUSTRIES		CFTYP_ALBVIS_SOIL	40
NAM_DATA_TEB	43	NAM_DATA_ISBA	40
CFNAM_H_TRAFFIC		CFTYP_ALBVIS_VEG	40
NAM_DATA_TEB	43	NAM_DATA_ISBA	42
CFNAM_H_TREE		CFTYP_BLD	42
NAM_DATA_ISBA	41	NAM_DATA_TEB	42
CFNAM_HC_ROAD		CFTYP_BLD_HEIGHT	

NAM_DATA_TEB	42	NAM_FRAC	37
CFTYP_BSLAI		CFTYP_SEFOLD	
NAM_DATA_ISBA	41	NAM_DATA_ISBA	41
CFTYP_CE_NITRO		CFTYP_SST	
NAM_DATA_ISBA	41	NAM_DATA_SEAFLUX	44
CFTYP_CF_NITRO		CFTYP_TC_ROAD	
NAM_DATA_ISBA	41	NAM_DATA_TEB	42
CFTYP_CNA_NITRO		CFTYP_TC_ROOF	
NAM_DATA_ISBA	41	NAM_DATA_TEB	42
CFTYP_CV		CFTYP_TC_WALL	
NAM_DATA_ISBA	40	NAM_DATA_TEB	43
CFTYP_D_ROAD		CFTYP_TOWN	
NAM_DATA_TEB	42	NAM_FRAC	37
CFTYP_D_ROOF		CFTYP_VEG	
NAM_DATA_TEB	42	NAM_DATA_ISBA	39
CFTYP_D_WALL		CFTYP_VEGTYPE	
NAM_DATA_TEB	43	NAM_DATA_ISBA	38
CFTYP_DG		CFTYP_WALL_O_HOR	
NAM_DATA_ISBA	39	NAM_DATA_TEB	42
CFTYP_DMAX		CFTYP_WATER	
NAM_DATA_ISBA	41	NAM_FRAC	37
CFTYP_EMIS		CFTYP_WRM_MAX_CF	
NAM_DATA_ISBA	39	NAM_DATA_ISBA	40
CFTYP_EMIS_ROAD		CFTYP_Z0	
NAM_DATA_TEB	42	NAM_DATA_ISBA	39
CFTYP_EMIS_ROOF		CFTYP_Z0_O_Z0H	
NAM_DATA_TEB	42	NAM_DATA_ISBA	40
CFTYP_EMIS_WALL		CFTYP_Z0_TOWN	
NAM_DATA_TEB	43	NAM_DATA_TEB	42
CFTYP_F2I		CGRID	
NAM_DATA_ISBA	41	NAM_PGD_GRID	47
CFTYP_GAMMA		CHORT	
NAM_DATA_ISBA	40	NAM_SGH_ISBA _n	70
CFTYP_GC		CISBA	
NAM_DATA_ISBA	41	NAM_ISBA	55
CFTYP_GMES		CKSAT	
NAM_DATA_ISBA	41	NAM_SGH_ISBA _n	70
CFTYP_H_INDUSTRIES		CLAMBERT	
NAM_DATA_TEB	43	NAM_IGN	51
CFTYP_H_TRAFFIC		CNATURE	
NAM_DATA_TEB	43	NAM_PGD_SCHEMES	46
CFTYP_H_TREE		COROGTYPE	
NAM_DATA_ISBA	41	NAM_ZS	54
CFTYP_HC_ROAD		CPHOTO	
NAM_DATA_TEB	42	NAM_ISBA	55
CFTYP_HC_ROOF		CRAIN	
NAM_DATA_TEB	42	NAM_SGH_ISBA _n	70
CFTYP_HC_WALL		CROUGH	
NAM_DATA_TEB	43	NAM_ISBA _n	70
CFTYP_LAI		CRUNOFF	
NAM_DATA_ISBA	39	NAM_SGH_ISBA _n	70
CFTYP_LAIMIN		CSCOND	
NAM_DATA_ISBA	41	NAM_ISBA _n	70
CFTYP_LE_INDUSTRIES		CSEA	
NAM_DATA_TEB	43	NAM_PGD_SCHEMES	46
CFTYP_LE_TRAFFIC		CSEA_ALB	
NAM_DATA_TEB	43	NAM_SEAFLUX _n	68
CFTYP_NATURE		CSEA_FLUX	
NAM_FRAC	37	NAM_SEAFLUX _n	68
CFTYP_RE25		CSNOW	
NAM_DATA_ISBA	41	NAM_PREP_ISBA_SNOW	64
CFTYP_RGL		CSNOWRES	
NAM_DATA_ISBA	40	NAM_ISBA _n	70
CFTYP_ROOTFRAC		CSOILFRZ	
NAM_DATA_ISBA	39	NAM_ISBA _n	70
CFTYP_RSMIN		CTOPREG	
NAM_DATA_ISBA	40	NAM_SGH_ISBA _n	70
CFTYP_SEA		CTOWN	

NAM_PGD_SCHEMES	46	NAM_PREP_SEAFLUX	61
CTYPE		LSSEDIMENTS	
NAM_PREP_FLAKE	63	NAM_FLAKE _n	69
NAM_PREP_ISBA	64	LSST_DATA	
NAM_PREP_SEAFLX	61	NAM_DATA_SEAFLUX	44
NAM_PREP_TEB	65	LSURF_BUDGET	
NAM_PREP_WATFLX	62	NAM_DIAG_SURF _n	74
CTYPE_TG		LSURF_BUDGETC	
NAM_PREP_ISBA	64	NAM_DIAG_ISBA _n	75
CTYPE_TS		LSURF_MISC_BUDGET	
NAM_PREP_TEB	65	NAM_DIAG_ISBA _n	75
CTYPE_WG		NAM_DIAG_TEB _n	75
NAM_PREP_ISBA	64	LSURF_VARS	
CTYPE_WS		NAM_DIAG_SURF _n	74
NAM_PREP_TEB	65	LTEB_CANOPY	
CWATER		NAM_PREP_TEB	65
NAM_PGD_SCHEMES	46	LWAT_SBL	
		NAM_PREP_SEAFLUX	62
I			
IDXRATIO		N	
NAM_INIFILE_CARTESIAN	49	N2M	
NAM_INIFILE_CONF_PROJ	48	NAM_DIAG_SURF _n	74
IDYRATIO		NDAY	
NAM_INIFILE_CARTESIAN	49	NAM_PREP_FLAKE	63
NAM_INIFILE_CONF_PROJ	48	NAM_PREP_ISBA	64
IXOR		NAM_PREP_SEAFLX	61
NAM_INIFILE_CARTESIAN	49	NAM_PREP_SURF_ATM	60
NAM_INIFILE_CONF_PROJ	48	NAM_PREP_TEB	65
IXSIZE		NAM_PREP_WATFLX	62
NAM_INIFILE_CARTESIAN	49	NDAY_SST	
NAM_INIFILE_CONF_PROJ	48	NAM_DATA_SEAFLUX	44
IYOR		NDGLG	
NAM_INIFILE_CARTESIAN	49	NAMDIM	52
NAM_INIFILE_CONF_PROJ	48	NDUMMY_PGD_NBR	
IYSIZE		NAM_DUMMY_PGD	56
NAM_INIFILE_CARTESIAN	49	NEMIS_PGD_NBR	
NAM_INIFILE_CONF_PROJ	48	NAM_CH_EMIS_PGD	57
		NEMIS_PGD_TIME	
L		NAM_CH_EMIS_PGD	57
LALDTHRES		NGROUND_LAYER	
NAM_SURF_ATM	67	NAM_ISBA	55
LALDZ0H		NIMAX	
NAM_SURF_ATM	67	NAM_CARTESIAN	49
LCH_BIO_FLUX		NAM_CONF_PROJ_GRID	48
NAM_CH_ISBA _n	72	NJMAX	
LCH_SURF_EMIS		NAM_CARTESIAN	49
NAM_CH_SURF _n	71	NAM_CONF_PROJ_GRID	48
LCOEF		NLAT	
NAM_DIAG_SURF _n	74	NAM_CARTESIAN	50
LFLK_SBL		NLON	
NAM_PREP_SEAFLUX	63	NAM_CARTESIAN	50
LFRAC		NMONTH	
NAM_DIAG_SURF_ATM _n	73	NAM_PREP_FLAKE	63
LISBA_CANOPY		NAM_PREP_ISBA	64
NAM_PREP_ISBA	64	NAM_PREP_SEAFLX	61
LPGD		NAM_PREP_SURF_ATM	60
NAM_DIAG_ISBA _n	75	NAM_PREP_TEB	65
LPRECIP		NAM_PREP_WATFLX	62
NAM_SEAFLUX _n	68	NMONTH_SST	
LPWEBB		NAM_DATA_SEAFLUX	44
NAM_SEAFLUX _n	68	NPATCH	
LPWG		NAM_ISBA	55
NAM_SEAFLUX _n	68	NPOINTS	
LRAD_BUDGET		NAM_IGN	51
NAM_DIAG_SURF _n	74	NRGRI	
LRM_TOWN		NAMRGRI	52
NAM_COVER	53	NTIME	
LSEA_SBL		NAM_DATA_ISBA	39

NAM_DATA_SEAFLUX	44	XLONMIN	
NYEAR		NAM_LONLMAT_REG	50
NAM_PREP_FLAKE	63	XRM_COVER	
NAM_PREP_ISBA	64	NAM_COVER	53
NAM_PREP_SEAFLX	61	XRPK	
NAM_PREP_SURF_ATM	60	NAM_CONF_PROJ	48
NAM_PREP_TEB	65	XSST_UNIF	
NAM_PREP_WATFLX	62	NAM_PREP_SEAFLX	61
NYEAR_SST		XTG_DEEP	
NAM_DATA_SEAFLUX	44	NAM_PREP_ISBA	64
NZSFILTER		XTG_ROOT	
NAM_ZS	54	NAM_PREP_ISBA	64
		XTG_SURF	
		NAM_PREP_ISBA	64
R		XTI_BLD	
RLOCEN		NAM_PREP_TEB	65
NAMGEM	52	XTI_ROAD	
RMUCEN		NAM_PREP_TEB	65
NAMGEM	52	XTIME	
RSTRET		NAM_PREP_FLAKE	63
NAMGEM	52	NAM_PREP_ISBA	64
		NAM_PREP_SEAFLX	61
		NAM_PREP_SURF_ATM	60
		NAM_PREP_TEB	65
		NAM_PREP_WATFLX	62
X		XTIME_SST	
XBETA		NAM_DATA_SEAFLUX	44
NAM_CONF_PROJ	48	XTS_ROAD	
XCGMAX		NAM_PREP_TEB	65
NAM_ISBA _n	70	XTS_ROOF	
XCISMIN		NAM_PREP_TEB	65
NAM_SURF_ATM	67	XTS_WALL	
XDX		NAM_PREP_TEB	65
NAM_CARTESIAN	49	XTS_WATER_UNIF	
NAM_CONF_PROJ_GRID	48	NAM_PREP_FLAKE	63
NAM_IGN	51	NAM_PREP_WATFLX	62
XDY		XTSTEP	
NAM_CARTESIAN	49	NAM_ISBA _n	70
NAM_CONF_PROJ_GRID	48	XUNIF_ALB_ROAD	
NAM_IGN	51	NAM_DATA_TEB	42
XEDB		XUNIF_ALB_ROOF	
NAM_SURF_ATM	67	NAM_DATA_TEB	42
XEDC		XUNIF_ALB_WALL	
NAM_SURF_ATM	67	NAM_DATA_TEB	43
XEDD		XUNIF_ALBNIR_SOIL	
NAM_SURF_ATM	67	NAM_DATA_ISBA	40
XEDK		XUNIF_ALBNIR_VEG	
NAM_SURF_ATM	67	NAM_DATA_ISBA	40
XENV		XUNIF_ALBUV_SOIL	
NAM_ZS	54	NAM_DATA_ISBA	40
XHUG_DEEP		XUNIF_ALBUV_VEG	
NAM_PREP_ISBA	64	NAM_DATA_ISBA	40
XHUG_ROOT		XUNIF_ALBVIS_SOIL	
NAM_PREP_ISBA	64	NAM_DATA_ISBA	40
XHUG_SURF		XUNIF_ALBVIS_VEG	
NAM_PREP_ISBA	64	NAM_DATA_ISBA	40
XLAT0		XUNIF_BLD	
NAM_CARTESIAN	49	NAM_DATA_TEB	42
NAM_CONF_PROJ	48	XUNIF_BLD_HEIGHT	
XLATCEN		NAM_DATA_TEB	42
NAM_CONF_PROJ_GRID	48	XUNIF_BSLAI	
XLATMAX		NAM_DATA_ISBA	41
NAM_LONLMAT_REG	50	XUNIF_CE_NITRO	
XLATMIN		NAM_DATA_ISBA	41
NAM_LONLMAT_REG	50	XUNIF_CF_NITRO	
XLON0		NAM_DATA_ISBA	41
NAM_CARTESIAN	49	XUNIF_CLAY	
NAM_CONF_PROJ	48	NAM_ISBA	55
XLONCEN			
NAM_CONF_PROJ_GRID	48		
XLONMAX			
NAM_LONLMAT_REG	50		

XUNIF_CNA_NITRO		XUNIF_ROOTFRAC	
NAM_DATA_ISBA	41	NAM_DATA_ISBA	39
XUNIF_COVER		XUNIF_RSMIN	
NAM_COVER	53	NAM_DATA_ISBA	40
XUNIF_CV		XUNIF_RUNOFFB	
NAM_DATA_ISBA	40	NAM_ISBA	55
XUNIF_D_ROAD		XUNIF_SAND	
NAM_DATA_TEB	42	NAM_ISBA	55
XUNIF_D_ROOF		XUNIF_SEA	
NAM_DATA_TEB	42	NAM_FRAC	37
XUNIF_D_WALL		XUNIF_SEFOLD	
NAM_DATA_TEB	43	NAM_DATA_ISBA	41
XUNIF_DEPTH_BS		XUNIF_T_B1	
NAM_DATA_FLAKE	45	NAM_PREP_FLAKE	63
XUNIF_DG		XUNIF_T_BOT	
NAM_DATA_ISBA	39	NAM_PREP_FLAKE	63
XUNIF_DMAX		XUNIF_T_BS	
NAM_DATA_ISBA	41	NAM_DATA_FLAKE	45
XUNIF_EMIS		XUNIF_T_ICE	
NAM_DATA_ISBA	39	NAM_PREP_FLAKE	63
XUNIF_EMIS_ROAD		XUNIF_T_MNW	
NAM_DATA_TEB	42	NAM_PREP_FLAKE	63
XUNIF_EMIS_ROOF		XUNIF_T_SNOW	
NAM_DATA_TEB	42	NAM_PREP_FLAKE	63
XUNIF_EMIS_WALL		XUNIF_TC_ROAD	
NAM_DATA_TEB	43	NAM_DATA_TEB	42
XUNIF_EXTCOEF_WATER		XUNIF_TC_ROOF	
NAM_DATA_FLAKE	45	NAM_DATA_TEB	42
XUNIF_F2I		XUNIF_TC_WALL	
NAM_DATA_ISBA	41	NAM_DATA_TEB	43
XUNIF_GAMMA		XUNIF_TOWN	
NAM_DATA_ISBA	40	NAM_FRAC	37
XUNIF_GC		XUNIF_VEG	
NAM_DATA_ISBA	41	NAM_DATA_ISBA	39
XUNIF_GMES		XUNIF_VEGTYPE	
NAM_DATA_ISBA	41	NAM_DATA_ISBA	38
XUNIF_H_B1		XUNIF_WALL_O_HOR	
NAM_PREP_FLAKE	63	NAM_DATA_TEB	42
XUNIF_H_ICE		XUNIF_WATER	
NAM_PREP_FLAKE	63	NAM_FRAC	37
XUNIF_H_INDUSTRIES		XUNIF_WATER_DEPTH	
NAM_DATA_TEB	43	NAM_DATA_FLAKE	45
XUNIF_H_SNOW		XUNIF_WATER_FETCH	
NAM_PREP_FLAKE	63	NAM_DATA_FLAKE	45
XUNIF_H_TRAFFIC		XUNIF_WRMAX_CF	
NAM_DATA_TEB	43	NAM_DATA_ISBA	40
XUNIF_H_TREE		XUNIF_Z0	
NAM_DATA_ISBA	41	NAM_DATA_ISBA	39
XUNIF_HC_ROAD		XUNIF_Z0_O_Z0H	
NAM_DATA_TEB	42	NAM_DATA_ISBA	40
XUNIF_HC_ROOF		XUNIF_Z0_TOWN	
NAM_DATA_TEB	42	NAM_DATA_TEB	42
XUNIF_HC_WALL		XUNIF_ZS	
NAM_DATA_TEB	43	NAM_ZS	54
XUNIF_LAI		XUSURIC	
NAM_DATA_ISBA	39	NAM_SURF_ATM	67
XUNIF_LAIMIN		XUSURICL	
NAM_DATA_ISBA	41	NAM_SURF_ATM	67
XUNIF_LE_INDUSTRIES		XUSURID	
NAM_DATA_TEB	43	NAM_SURF_ATM	67
XUNIF_LE_TRAFFIC		XVCHRNK	
NAM_DATA_TEB	43	NAM_SURF_ATM	67
XUNIF_NATURE		XVMODMIN	
NAM_FRAC	37	NAM_SURF_ATM	67
XUNIF_RE25		XVZ0CM	
NAM_DATA_ISBA	41	NAM_SURF_ATM	67
XUNIF_RGL		XWS_ROAD	
NAM_DATA_ISBA	40	NAM_PREP_TEB	65

XWS_ROOF	
NAM_PREP_TEB	65
XX	
NAM_IGN	51
Y	
YCLAY	
NAM_ISBA	55
YCLAYFILETYPE	
NAM_ISBA	55
YCOVER	
NAM_COVER	53
YDEPTH_BS	
NAM_DATA_FLAKE	45
YDEPTH_BSFILETYPE	
NAM_DATA_FLAKE	45
YEXTCOEF_WATER	
NAM_DATA_FLAKE	45
YEXTCOEF_WATERFILETYPE	
NAM_DATA_FLAKE	45
YFILETYPE	
NAM_COVER	53
NAM_PGD_GRID	47
NAM_ZS	54
YINIFILE	
NAM_PGD_GRID	47
YSAND	
NAM_ISBA	55
YSANDFILETYPE	
NAM_ISBA	55
YT_BS	
NAM_DATA_FLAKE	45
YT_BSFILETYPE	
NAM_DATA_FLAKE	45
YWATER_DEPTH	
NAM_DATA_FLAKE	45
YWATER_DEPTHFILETYPE	
NAM_DATA_FLAKE	45
YWATER_FETCH	
NAM_DATA_FLAKE	45
YWATER_FETCHFILETYPE	
NAM_DATA_FLAKE	45
YY	
NAM_IGN	51
YZS	
NAM_ZS	54